



Management of Multidrug-Resistant Tuberculosis in Children: **A FIELD GUIDE**

Fourth Edition: February 2019
Cover photo courtesy of Marcela Tommasi

This handbook was developed and written by The Sentinel Project for Pediatric Drug-Resistant Tuberculosis (<http://sentinel-project.org/>).

Preferred citation: Management of Drug-Resistant Tuberculosis in Children: A Field Guide. Boston, USA: The Sentinel Project for Pediatric Drug-Resistant Tuberculosis; November 2018, Fourth edition



The Sentinel Project on Pediatric Drug-Resistant Tuberculosis is a global partnership of researchers, caregivers, and advocates aiming to develop and deploy evidence-based strategies to prevent child deaths from this treatable disease. We are a learning network committed to generating and disseminating knowledge and data for immediate action.

The fourth edition of this handbook is based on the Management of Multidrug-Resistant Tuberculosis in Children: A Field Guide, First Edition, November 2012, Second Edition, March 2015, and Third Edition, November 2016 a resource developed by the Sentinel Project on Pediatric Drug- Resistant Tuberculosis.

Acknowledgments

The main writing committee for this handbook was: Jennifer Furin, James Seddon and Mercedes Becerra.

Other contributors included: Jay Achar, Farhana Amanullah, Annemieke Brands, Grania Brigden, Gunta Dravniece, Anthony Garcia-Prats, Malgorzata Grzemska, Anneke Hesseling, Kerry Holloway, Zheng Huiwen, Alexander Kay, Uzma Khan, Iqbal Master, Ben Marais, Lindsay McKenna, Carlos Perez-Velez, Anja Reuter, H. Simon Schaaf, Jeffrey Starke, Marcela Tommasi, Pilar Ustero, Sabine Verkuijl, Jessica Vineberg, Askar Yedilbayev, and Courtney Yuen.

Editorial comments on the Second Edition were provided by Agnes Gebhard, Maria Insua, Ieva Leimane, and Kwonjune Seung.

The following individuals contributed to earlier editions of this field guide that formed the basis of this fourth edition: Tonya Arscott-Mills, Karen Day, Hernán Del Castillo, Peter Drobac, Marianne Gale, Chelsie GawneMark, Elizabeth Haraus, J. Keith Joseph, Marina Marinkovic, , Sharon Nachman, Michael Rich, Uruvashi Singh, Soumya Swaminathan, Arielle Tolman, Zheng Yang, Zhao Yanlin, Askar Yedilbayev, and Courtney Yuen.

The authors of this handbook are thankful to countless people for the time, expertise, and generosity which led to the earlier editions of the handbook. First and foremost, we are mindful of the children and their families who face this disease on a daily basis. Their bravery inspires us, and it is for them that we aspire to zero childhood TB disease and death. For funding and moral support, we thank the Stop TB Partnership and the incredible individuals who work there. Finally, we thank the men and women who care for sick children around the world. May we all come to know a day when all children have the hope of happy and healthy lives.

The Fourth Edition of this Field Handbook is dedicated to **Ms. Shreya Tripathy**, a brave young woman whose battle with drug-resistant TB has been an inspiration to us all.

TABLE OF CONTENTS

INTRODUCTION	1
SECTION 1: DIAGNOSIS	2
Algorithm for Suspected MDR-TB	3
Specimen Management	7
SECTION 2: REGIMEN DESIGN	11
Principles	11
Dosing	21
SECTION 3: MONITORING	31
Schedule of Visits	31
Adverse Events.....	35
Co-morbid Conditions.....	39
Nutritional Assessment and Support	41
Adherence	43
Special Populations: Neonates and Adolescents.....	46
SECTION 4: MDR-TB CONTACTS	47
SECTION 5: INFECTION CONTROL	49
Facility-based.....	50
Community-based	51
SELECTED REFERENCES	53
APPENDICES	56
Appendix A: Sample Intake Form	56
Appendix B: Medications Used to Treat MDR-TB	61
Appendix C: Contact Management Form	63
Appendix D: Specimen Collection Procedures	65

Disclaimer

As between The Sentinel Project on Pediatric Drug-Resistant Tuberculosis (hereafter referred to as “The Sentinel Project”) and you, The Sentinel Project or its licensors at all times owns and retains all right, title, and interest in and to the “Management of Drug-Resistant Tuberculosis in Children: A Field Guide” including all intellectual property rights therein and thereto. You may use and copy the Field Guide,

or portions of the Field Guide, provided that you (i) promptly notify The Sentinel Project via email at sentinel_project@hms.harvard.edu of any such use and (ii) reproduce all copyright notices, claims, or reservation of rights appearing on the Field Guide, as delivered to you, on all copies made pursuant to this sentence.

By using the Field Guide, you acknowledge that (i) the Contributors and Authors are not guaranteeing the completeness or accuracy of the information contained in the Field Guide, (ii) the Field Guide does not represent nor comprise all of the data or other information that may be relevant to you, (iii) portions of the Field Guide may be outdated and may have been superseded by subsequent changes in applicable law, regulation, or conditions affecting nonprofit health organizations generally, and (iv) the Contributors and Authors do not have any duty to update the Field Guide or, if updated, to provide further updated versions of the Field Guide.

To the maximum extent permitted under applicable laws, none of the Contributors and Authors are liable, under any circumstance or under any theory of law, tort, contract, strict liability or otherwise, for any direct, indirect, special, incidental, consequential, exemplary or punitive damages, or any other injury and/or damage (actual or perceived), including, without limitation, damages based on errors, omissions, or reliance upon, use, or operation of any ideas, instructions, procedures, products, or methods contained in the material included in this guide.

By using this guide, you acknowledge and agree to the terms of this notice.



Photo courtesy of Jennifer Furin

INTRODUCTION

Multidrug-resistant (MDR) tuberculosis (TB) is a growing global health crisis; MDR-TB is defined TB disease caused by strains of *Mycobacterium tuberculosis* with in vitro resistance to at least isoniazid and rifampicin, and it is estimated there are more than five million people infected and sick with drug-resistant forms of TB in the world today. With the increasing use of the Xpert MTB/RIF® to detect both TB and rifampicin resistance, the term “rifampicin-resistant TB” (RR-TB) is increasingly used as well. In general, RR-TB is treated the same as MDR-TB and thus the term MDR-TB will be used in this field guide to encompass RR-TB as well.

Children represent a significant proportion of persons with TB disease with an estimated 30,000 children becoming sick with MDR-TB each year. Yet they lack the same access to diagnosis and treatment as their adult counterparts. Two meta-analyses of treatment for MDR-TB among children showed that about 80% have positive outcomes when treated for MDR-TB. These reviews also demonstrated that, with the exception of the injectable agents, children tolerated second-line medications well. However, fewer than 5% of children who become sick with MDR-TB are ever started on appropriate treatment for their disease.

Urgent action is needed to address this significant gap in care. Based on experiences with pediatric HIV, equitable access for children with MDR-TB will only occur once systematic approaches to diagnosing and treating children are developed and once access to pediatric formulations of second-line medications is widespread. This field guide is intended to serve as a tool for practitioners working with children at risk of MDR-TB infection and those who

are sick with MDR-TB disease. Following the example set by Médecins Sans Frontières (MSF) in their publication “Treating drug-resistant tuberculosis: what does it take,” this guide focuses on issues relevant in clinical and programmatic practice and does not offer extensive background materials on management of MDR-TB, which can be found in multiple other guidelines. This field guide should be considered complementary to existing recommendations.

In general, a guide such as this should be developed using evidence-based research. The World Health Organization (WHO) has established methodologies for the development of their guidelines that follow a systematic evaluation of available evidence. And although there are multiple planned and ongoing clinical trials focused on optimal treatment of MDR-TB infection and disease in children, to date, there are few completed studies to form the basis of pediatric-specific treatment guidelines. While many aspects of the adult recommendations apply to children, there are some unique aspects of pediatric MDR-TB that may require different approaches. While awaiting the results of pediatric clinical trials, however, practitioners in most countries are already seeing children at risk for, or sick with, MDR-TB, who require immediate access to high-quality care. This field guide was developed by a team of experts who jointly have treated thousands of children with MDR-TB over the last two decades in every region of the world. It combines the best available research evidence with clinical experience. We hope it will be used as a tool to rapidly increase the number of children receiving care for MDR-TB. Case examples are included throughout the guide to show how these recommendations can be put into practice. The WHO have been very supportive of the development of this field guide to provide advice that extends beyond their guidelines.

Whenever possible, management of children with MDR-TB should take place within the activities of a National TB Program (NTP). There are multiple advantages to doing this, including a contextual approach, integration with other health initiatives, and health systems strengthening. If activities occur outside of an NTP, all efforts should be made to report standardized outcomes and to collaborate with the NTP whenever possible.

Finally, we recognize that the term “children” encompasses a broad range of individuals and ages with widely different needs. A 2-year-old child requires a different approach to a 12-year-old, and the treatment of children with MDR-TB will never be a “one size fits all” approach. Many experts feel that children older than 10 years of age can be managed as adults using the same diagnostic methods and medications, although the specific emotional needs of adolescents and their caregivers should be considered. We focus mainly on the care of younger children, which is most problematic for most practitioners and hope to offer a compendium of practical experience that can be useful for programs and providers caring for children with MDR-TB. We trust that this initial attempt will be greatly expanded and improved as the world gains and documents more experience with this neglected population in the coming years.

1. DIAGNOSIS OF MDR-TB DISEASE IN CHILDREN

Key points to consider in MDR-TB diagnosis in children are:

- The diagnosis of MDR-TB in children is mostly made on clinical and radiological grounds with consideration of risk factors for MDR-TB (e.g. recent MDR-TB exposure)

- Bacteriological confirmation should be attempted, but is often not possible due to paucibacillary disease or extrapulmonary (EP) disease
- Bacteriological results are frequently negative even if the child has TB (sputum smear results in <15%, Xpert MTB/RIF or culture in ~40%)
- The collection quality respiratory samples can be difficult in young children, but children >5 years old can usually provide sputum. In younger children gastric aspiration/lavage or induced sputum can be done
- Other tests to diagnose EP-TB should be performed as needed

This section will discuss:

- Diagnosis of pediatric MDR-TB
- Diagnostic specimens, preparation, and testing

Recognition and Initial Management of a Child with Possible MDR-TB

TB should be included in the differential diagnosis list of any child with a persistent non-remitting cough or fever, weight loss/failure to thrive, or focal findings that are suggestive of TB, such as lymphadenitis, spinal deformities, ascites, and joint effusions. Danger signs of possible meningitis include lethargy/sleepiness, loss of consciousness, and seizures. Diagnosis of MDR-TB among children is can be challenging and requires a high level of suspicion. MDR-TB in children can either be confirmed (they have clinical TB disease and a sample taken from the child shows MDR-TB) or clinically diagnosed (the child has clinical TB disease and has risk factors for drug resistance). Clinically diagnosed TB includes probable and possible MDR-TB. Under field conditions, it may take several weeks from the time a child first presents with signs and symptoms of TB and the receipt of test results, during which time a child can rapidly deteriorate. Thus, it is important to consider initiating MDR-TB therapy in the absence of bacteriologic confirmation.

Some definitions to consider are:

- **Confirmed MDR-TB:** MDR-TB is isolated from the child
- **Probable MDR-TB:** Symptoms/ signs and/or radiology consistent with TB disease in a child who has been exposed to an adult with infectious MDR-TB (>80% concordance between drug susceptibility test (DST) patterns in diseased children and the likely source case)
- **Possible MDR-TB:** Child is not improving after 2-3 months of first-line treatment (with confirmation of treatment adherence and exclusion of likely alternative diagnosis)
or
Close contact with a patient who: died from TB; failed TB treatment or is a TB retreatment case

Figure 1 provides an approach for the investigation and management of possible and probable MDR-TB.

Figure 1: Approach to the management of possible and probable MDR-TB disease




Table 1: Abbreviations for drugs used in this Field Guide

Drug Name	Abbreviation
Amikacin	Am
Amoxicillin-clavulanic acid	Amx-Clv
Bedaquiline	Bdq
Capreomycin	Cm
Clofazimine	Cfz
Cycloserine	Cs
Delamanid	Dlm
Ethambutol	E
Ethionamide	Eto
Imipenem-cilastatin	Imp-Cln
Isoniazid	H
Kanamycin	Km
Levofloxacin	Lfx
Linezolid	Lzd
Meropenem	Mpm
Moxifloxacin	Mfx
Para-aminosalicylic acid	PAS
Prothionamide	Pto
Pyrazinamide	Z
Rifampicin	R
Streptomycin	S
Terizidone	Trd

Note: In the Field Guide, to avoid confusion we write out drugs in full when describing the individual drug. Where we describe regimens, we use the abbreviations above.

Case Examples: Possible MDR-TB in a Child

Colombia

Antonio is a 7-year-old boy who presents to the clinic with 3 weeks of fever, cough, and a swollen right knee. His father notes that the boy has been “sleepy” at school and is no longer interested in playing football with his friends. When asked how he is doing, Antonio says his “leg hurts” but reports no other complaints. His father also tells you that Antonio’s mother died last month. “They say it was from TB, but I do not know.

She took her TB medication every single day with the nurse watching her.” There are no other known TB exposures, and prior to this episode, Antonio has been healthy, growing well, and has received all his recommended vaccinations, including BCG vaccination which is verified by the presence of a scar.

On examination, Antonio is pale and listless. His temperature is 38.1°C and his weight is only 16 kg, which is below the fifth percentile for him and a drop from his growth

curve, where he had previously been at the fifteenth percentile. He has cervical lymphadenopathy and examination of the right knee shows arthritis and a small effusion.

You suspect TB and are concerned about the possibility of MDR-TB, given that his mother died of TB while on first-line treatment even though she had excellent adherence. Antonio is clinically stable with no immediate indication to start MDR-TB treatment. He provides sputum for smear and culture, and a sample of his knee fluid is sent to the lab for analysis as well. You inquire about the use of a rapid genotypic test (GeneXpert) at the National Lab. His chest radiograph shows a patchy right upper lobe infiltrate. All sample results are negative when his father brings him back to the clinic with a temperature of 40.5°C and in respiratory distress. Given his clinical instability and risk factors for MDR-TB, you start him on an MDR-TB regimen in the absence of bacteriologic confirmation. MDR-TB is never confirmed. He does well with this treatment and is cured of his presumed MDR-TB.

Kenya

Shamba is a 7-year-old boy who is brought to the health center by his mother after she notices “lumps” in his neck and that the child is “coughing all the time.” Shamba says he feels tired and “sweaty all the time.” He begins to cry and tells you he misses his father, who died last year. Upon further questioning, Shamba’s mother reports in a whisper that her husband died of TB last year, even though he took his TB medications every day.

On examination, Shamba is noted to be cachectic. He is febrile to 39°C and tachycardic with a heart rate of 137. On exam, he has multiple cervical lymph nodes, which are cool to the touch but feel round and rubbery. His cardiac exam reveals a 2/6 systolic ejection murmur. His pulmonary exam reveals chest wall retractions with minimal expansion of the right hemithorax. The right side of the chest is dull to percussion, and no breath sounds can be heard.

Shamba is critically ill and undergoes an emergency thoracentesis with removal of 600 ml of straw-colored fluid. He is able to give a sputum sample, which is sent for smear, culture, and rapid DST. Because he is so ill and given the fact that his father died of TB while in treatment, Shamba is started on treatment for MDR-TB. Two days later, the results from his Xpert MTB/RIF™ show resistance to rifampicin, and his final culture comes back with resistance to isoniazid, rifampicin, ethambutol and streptomycin. Shamba is able to fully recover and remains an active, playful 9-year-old (having completed 12 months of treatment when he was seven) at the top of his class in school.

Kazakhstan

Aizhan is an 11-year-old girl who presents to her health center with fever, weight loss, cough, and hemoptysis. Aizhan is in her fourth month of TB treatment and is currently on the continuation phase of treatment given through her National TB Program using DOTS.

Aizhan was originally diagnosed 4 months ago when she presented with the same symptoms and was found to have a positive acid-fast bacilli (AFB) smear: unfortunately, no Xpert or culture testing was done. She was started on HRZE and received daily supervised therapy at a TB dispensary in her community. She has not missed a single dose of therapy. In her first month of treatment, she had a negative smear and reported feeling “better” with resolution of her cough and fever. During her second month

of treatment, however, she began to cough again and developed drenching night sweats. Her month two sputum smear was negative for AFB, and she was put on the continuation phase (isoniazid and rifampicin) after the second month of therapy. Her symptoms worsened, and she began to have daily fevers. She presented to her clinic, and her month three smear showed “rare AFB,” which was felt to be a “contamination” by her providers when a repeat smear was negative. She continued on isoniazid and rifampicin. Her daily compliance with therapy was confirmed.

She now presents to the health center after coughing blood-streaked sputum. She also notes a 6 kg weight loss, daily fevers, severe cough, and shortness of breath. On exam she is ill-appearing, cachectic, and tachycardic, and her lungs have diffuse crackles and wheeze throughout all fields. Her sputum is streaked with blood, but a smear is done which shows AFB. A repeat history is taken, and Aizhan and her mother deny any contacts with other TB patients and specifically state that they have no known contact with persons with MDR-TB or other risk factors. A rapid HIV test is negative.

Aizhan is deemed to be at high risk for MDR-TB, as she is failing a first-line regimen despite excellent adherence. Her sputum is sent for culture and DST. In the meantime, she is started on a regimen for her presumptive MDR-TB. The DST returns and shows resistance to isoniazid, rifampicin, ethambutol and streptomycin.

KEY POINTS:

- A high level of clinical suspicion is needed for timely diagnosis of MDR-TB in children.
- If the child has symptoms/signs/radiology of TB and has been exposed to an infectious MDR-TB case, then they should be considered to have probable MDR-TB
- Risk factors for possible MDR-TB include a history of previous treatment (especially within the past 12 months), failure to improve clinically on first-line TB treatment after 2 to 3 months of therapy (if IRIS not considered), contact with a patient who died on TB treatment or failed TB treatment.
- Treatment in the absence of bacteriologic confirmation should be commonly carried out
- In the case of probable MDR-TB, treatment should be based on the DST of the source case
- Early initiation of appropriate treatment is essential to ensure good outcomes.

(Photo credit Marcela Tommasi)



Diagnostic Specimen Management

There are multiple specimen types that can be taken from children to diagnose MDR-TB, and these can be sent for a variety of tests, including smear, liquid medium culture (i.e.

MGIT), solid medium culture, pathology, or rapid diagnostic testing with the GeneXpert® Ultra, GeneXpert®, or a line probe assay (i.e. GenoType® MTBDRplus). Of note, GeneXpert® Ultra and culture in liquid media should be prioritized in children. All relevant and available tests should be considered; performing multiple tests on one or more samples of a variety of specimen types significantly increases the diagnostic yield. The principles of specimen collection and management are described below. See Appendix A for a sample specimen collection form.

Infection Control Precautions

The proper collection of a specimen for microbiological studies (especially Xpert MTB/RIF® and cultures) is the most important step in the detection of *M. tuberculosis*. A poorly collected specimen may lead to failure in detecting *M. tuberculosis* and/or result in the recovery of contaminating organisms (including non-tuberculous mycobacteria). Whenever possible, specimens should be taken prior to starting TB treatment. For proper specimen collection:

- Collect the specimen at optimal times (e.g. early morning fasting gastric aspirate, before mobilization; induced sputum after fasting 2-4 hours; expectorated sputum early morning).
- Always try to collect an optimal quantity of sample, which varies by specimen type; larger volumes generally provide higher bacteriological yields.
- Use appropriate collection devices (i.e., sterile, leak-proof specimen containers). If specimens (e.g., sputa) will require centrifugation, they should preferably be collected directly into 50-mL centrifuge (Falcon) tubes to avoid the need for their transfer from one container to another.
- Avoid contamination, by only sampling the secretions/tissues of interest and adhering to basic sterile/clean procedure principles
- Use appropriate transport media. Avoid saline or other solutions that may contain antimycobacterial preservatives, do not place biopsy samples in formaldehyde, and neutralize stomach acid if gastric aspirate is collected (see Appendix D).
- If possible, collect specimens prior to administration of anti-tuberculosis medications. In critically ill children, this may not be possible, but all attempts should be made to obtain samples as quickly as possible.
- Properly label each specimen (complete names; exact type of specimen source; date & time collected). Ensure that the label is on the container, not the lid.
- Minimize transport time.
- Maintain an appropriate environment (cool temperature, ideally between 5-15°C) between collection of specimens and delivery to the laboratory.
- If a delay in transport is anticipated place specimen in refrigerator (ideally between 1-5°C); freeze specimens for molecular (PCR) analysis.
- Package each specimen separately in a sealed transport bag.

Standard Operating Procedures for the most relevant samples are supplied in Appendix D.

Types of Specimens

(All of these samples can be tested on Xpert and in culture, with the exception of pleural fluid for which Xpert is not recommended)

Table 2. Types of respiratory specimen collection and their conditions of use

Specimen	Brief description of sample collection procedure	Recommended age group	Recommended minimum volume for studies*	Optimal collection time	Comments/ tips
Spontaneous expectorated sputum (ES) expectorated sputum (ES)	Expectoration of sputum without prior saline nebulization	>5 years	3 mL	Early morning	If child is unable to produce sputum of sufficient quantity and quality, consider sputum induction.
Induced sputum (IS)	Expectoration of sputum preceded by hypertonic saline nebulization	Any age	3 mL	Early morning	If child is unable to expectorate (children under 5 years), consider laryngo-pharyngeal suctioning.
Gastric aspirate (GA)	Nasogastric aspiration of gastric juice containing swallowed sputum	<7 years	5 mL	Early morning before child gets out of bed	Upon awakening and sitting and standing, peristalsis begins, and stomach gradually empties, consequently compromising volume.
Gastric lavage (GL)	Nasogastric instillation of solution to “wash off” and recover sputum adhered to walls of stomach	<7 years	10 mL	Early morning	Recommended only when 3 mL of gastric aspirate cannot be obtained.
Broncho-alveolar lavage (BAL)	Bronchoscopy	Any age	3 mL	Any time	Bacteriologic yield of one BAL sample is not superior to that of serial induced sputum or GA/GL.
Nasopharyngeal aspirate	Nasopharyngeal suctioning of the nasopharynx to collect secretions from URT, but may also collect secretions from LRT if cough reflex is stimulated	<7 years	2 mL	Unknown, but probably higher yield in morning	The bacteriologic yield of naso-pharyngeal aspirate tends to be similar to, or lower than, that of induced sputum or GA/GL and this may be a good alternative to these methods of obtaining sputum.

* These values are the minimum recommended amount; larger volumes tend to have higher bacteriological yields.

Table 3: Types of non-respiratory specimen collection and their conditions of use

Specimen	Brief description of sample collection procedure	Recommended age group	Recommended minimum volume for studies*	Optimal collection time	Comments/ tips
Stool	Sampling of random stool uncontaminated by urine or toilet bowl	Any age	1 table-spoon (5 g)	Any time	Bacteriologic yield of stool for smear and culture has so far been lower than that of sputum and GA/GL. However, there are important and emerging data to show that Xpert testing done on stool after simple decontamination procedures can have a diagnostic yield similar to gastric aspirate and is much less invasive.
Cerebrospinal fluid (CSF)	Lumbar puncture	Any age	2 mL	Any time	Submit the 3rd or 4th tube for culture to reduce the possibility of contamination due to skin flora.
Serosal fluids and tissues**	Serosal fluid aspirate followed by serosal tissue biopsy	Any age	1 mL	Any time	Bacteriologic yield of tissue is significantly higher than that of fluid. Biochemical markers useful in all fluids.
Urine	Clean-catch, midstream urine	Any age	2 mL	First micturition of the morning	Bacteriologic yield low except in urinary tract TB. Detection of lipoarabinomannan antigen has high sensitivity in severely immuno- compromised HIV- infected patients. Of note, there is increasing evidence that the use of the urinary lipoarabinomannan (LAM) may be a useful test to diagnose TB in children or individuals living with HIV with low CD4 counts

Specimen	Brief description of sample collection procedure	Recommended age group	Recommended minimum volume for studies*	Optimal collection time	Comments/ tips
Blood	Phlebotomy	Any age	5 mL	Any time	Bacteriologic yield very low; use in severely ill HIV-infected patients.
Fine needle aspiration biopsy	Fine needle aspiration biopsy, depending on type of tissue and clinical situation	Any age	Based on type	Any time	Also useful because histopathologic features consistent with TB can be diagnostic. Note that this data refers to the biopsy only and not to a fine needle aspirate of fluid.
Bone marrow	Bone marrow aspirate	Any age	1mL	Any time	Consider bone marrow aspiration in children with disseminated disease. Should test for other pathogens, especially in children with HIV.

** Serosal fluids include pleura, pericardium, peritoneum, and synovium.

2. REGIMEN DESIGN

The medications used for the treatment of children with MDR-TB are similar to those used to treat adults. However, many second-line drug formulations are not child-friendly, and preparation can be labor intensive. However, there are now child-friendly, quality assured formulations of the following medications that can be obtained from the Global Drug Facility: pyrazinamide, ethambutol, levofloxacin, moxifloxacin, ethionamide, isoniazid, and cycloserine.

(http://www.stoptb.org/gdf/drugsupply/drugs_available.asp).

- Pharmacokinetic and pharmacodynamic testing of second-line TB drugs is increasingly being done in children and there are now data to support many of the recommended doses of the second-line drugs. This section will discuss:
- Principles of regimen design in children
- Dosing recommendations for second-line TB drugs in children

Pictures of the commonly used first- and second-line drugs are included in Appendix B.

Principles of Regimen Design: Pediatric Considerations

In 2018 the World Health Organization updated their guidance on the management of RR/MDR-TB. Chief among these recommendations were a re-prioritizing of the second-line

drugs into Group A, Group B and Group C medications. Although the data supporting this drug re-classification largely came from adults, the principles of regimen design also apply for children. A summary of the new drug groupings follows below:

Table 4. The new 2018 drug grouping for the treatment of MDR-TB

Group	Drug	Abbreviation
Group A Include all three drugs (unless they cannot be used)	Levofloxacin OR Moxifloxacin	Lfx Mfx
	Bedaquiline	Bdq
	Linezolid	Lzd
Group B Add both drugs (unless they cannot be used)	Clofazimine	Cfz
	Cycloserine OR Terizidone	Cs Trd
Group C Add to complete regimen and when drugs from Groups A and B cannot be used	Ethambutol	E
	Delamanid	Dlm
	Pyrazinamide	Z
	Imipenem-cilastatin	lpm-Cln
	Meropenem	Mpm
	Amikacin OR Streptomycin	Am S
	Ethionamide OR Prothionamide	Eto Pto
	p-aminosalicylic acid	PAS

In general, children with MDR-TB should be managed according to the same principles that guide adult therapy. For children, however, the following principles are recommended based on the consensus opinion of the experts involved in writing this field guide:

- Treatment should be based on the DST pattern of the most likely source case if the child does not have a DST of his or her own;
- Always attempt to treat children with injectable-free regimens, especially very young children and those with mild disease. Hearing loss, a frequent severe adverse event caused by injectables can have a profound impact on language acquisition and ability to learn at school and further development. This treatment principle is consistent with the 2018 WHO recommendations which state: ““The avoidance of an injectable-containing regimen is particularly desirable in children, especially those very young and with mild disease, as determined by the absence of malnutrition, serious forms of extrapulmonary disease, cavitation on chest radiography or HIV infection. Given the profound impact that hearing loss can have on the acquisition of language and the ability to learn at school, the use of amikacin or streptomycin in children is to be accompanied with regular audiometry.”
- Recommendations on MDR-TB treatment regimens for adults also apply to children with severe forms of extra-pulmonary MDR-TB. Treatment of MDR-TB meningitis should be guided by the medicines’ ability to cross the blood-brain barrier. See Table 7 for more details.

- Regimens should consist of at least 4 drugs to which the organism is likely to be susceptible for the duration of therapy, with possible addition of a 5th drug for the first few months of therapy in cases with severe or multi-bacillary. Using more than 5 drugs adds to the toxicity without necessarily improving treatment efficacy, if Group A and B drugs and/or delamanid are used;
- Regimen construction should prioritize the WHO Group A and B drugs, as well as delamanid in children aged more than 3 years of age (see Figure 2). PAS is an alternative drug to use in children instead of injectable agents or in instances where the new drugs are not yet available, although its association with worse outcomes in adults should be considered
- The WHO recommends bedaquiline for the treatment of children aged 6 years and above and the use of delamanid for the treatment of children aged 3 years and above. These newer drugs, bedaquiline and delamanid, have not been studied extensively in very young children, therefore, designing regimens for these children can be tricky. Although regimens will need to be designed for each individual patient—taking into account unique resistance patterns and toxicity risks, Figure 3 can be used to guide the design of treatment regimens for children with MDR-TB. It should be noted that although licensing for both delamanid and bedaquiline is usually for six months, there are no known safety concerns with using these drugs for longer than six months (although data are limited). Some children may benefit from using these drugs for the full duration of their therapy. Furthermore, there are limited data on the use of bedaquiline and delamanid in combination—although existing data suggest the combination of the two medications does not result in any increase in adverse events. In children with fluoroquinolone resistance or in whom there are limited treatment options, extension and combination of bedaquiline and/or delamanid could be considered on a patient-by-patient basis with careful monitoring. More data on the use of bedaquiline and delamanid in children are expected in the coming years and will be used to further refine these management principles.
- Although linezolid is a Group A drug with proven effectiveness, its use has been associated with frequent toxicity. Toxicity is duration dependent and although use throughout treatment is likely to improve efficacy, adverse events may limit the duration of use to the first few months
- Amoxicillin-Clavulanic acid should be administered with every dose of imipenem-cilastatin or meropenem to aid its efficacy. It should not be counted as an additional medicine or used as a separate agent
- Pyrazinamide should only be used if there is demonstrated susceptibility
- The composition of MDR-TB treatment regimens is largely the same in children living with HIV. Efavirenz should be avoided in children who need bedaquiline for the duration of their bedaquiline treatment, as efavirenz lowers the concentrations of bedaquiline.
- The duration of therapy in children should depend upon the site and severity of disease: children with non-severe disease can be treated for 9 to 12 months while children with severe disease will require 12-18 months of therapy depending on their clinical progress (see Figure 4). Of note, the 2018 WHO recommendations define

severe disease as follows: “In children <15 years, severe disease is usually defined by the presence of cavities or bilateral disease on chest radiography or extrapulmonary forms of disease other than lymphadenopathy (peripheral nodes or isolated mediastinal mass without compression). In children the occurrence of advanced malnutrition (defined by syndrome or by metrics) or advanced immunosuppression or positive TB bacteriology (smear, Xpert MTB/RIF, culture) may also be considered when determining treatment duration.

- Child-friendly formulations of the medications should be used whenever possible.
- Monitoring and management of adverse events is essential;

TB occurs along a spectrum ranging from infection to mild disease to severe disease, and the severity of disease is often related to bacillary burden in children. Because of this, children with non-severe disease can usually be treated with shorter durations of therapy. Non-severe disease can be defined as TB disease that is isolated to the lymph nodes or only affects one of the lungs without cavities. In general children with non-severe disease only need 9-12 months of treatment. For most children, treatment lasting longer than 18 months will not be necessary, and clinical experience suggests shorter durations of treatment (i.e. 15 to 18 months) could be considered in children with severe disease based on their clinical evolution, their comorbidities, and their DST patterns.

These principles are the same as for adults with MDR-TB and should be followed as far as possible. As there are less data on the use of delamanid in children under 3 years and bedaquiline in children under 6 years, we have constructed a series of possible regimens for children <3 years, 3 to <6 years and older than 6 years. For each age group we have suggested regimens to treat isolates that are either susceptible or resistant to the fluoroquinolones (Figure 3).

Figure 2. Principles in the construction of an MDR-TB regimen for children





Figure 3. Suggested regimens for children of different ages and with different resistant profiles to the fluoroquinolones



Rationale for recommendations in Figure 3

Figure 3 outlines a possible approach to regimen design based on different age groups and the presence or absence of fluoroquinolone resistance. These proposed regimens reflect the reality of regimen design in children, where there are limited data and where clinical experience still plays a major role in regimen construction. In this figure, additional drugs are suggested, based on clinical practice and could be added to strengthen a regimen (in case of severe disease) or could be used as a substitute if there is toxicity or intolerance to other drugs.

The age groups were selected based on the presence of pharmacokinetic and safety data to support the use of the medications. For example, WHO now recommends that bedaquiline can be given to children ages 6 years and above, and in this age group, bedaquiline is prioritized in treatment regimens. WHO also recommends that delamanid can be given to


children aged 3 years and above, and in this age group, delamanid is prioritized in regimen construction. Because of this, we have chosen to use the age bands 6 years and older, 3 to <6 years, under 3 years.

For children who are age 6 years and over and who have fluoroquinolone susceptible TB, the core of the regimen should be Bdq-Lfx-Lzd-Cfz. If a fifth drug is needed, then a choice can be made between cycloserine (a Group B drug with potential for neurological toxicity) or delamanid (a Group C drug that is relatively safe but for which there is limited safety data when used in combination with bedaquiline). Of note, ethambutol could be given if there is documented susceptibility to it and the same with pyrazinamide. For children with fluoroquinolone resistance, the risk of a poor outcome is increased and thus the use of both cycloserine and delamanid is justified. Some treating centers, however, report good clinical experience using PAS, and this drug could be considered for children with fluoroquinolone-resistant disease. PAS was associated with worse treatment outcomes in adults, however, and this should be taken into consideration in regimen design.

For children who are 3 to <6 years of age, the use of bedaquiline is not routinely recommended since there are not yet pharmacokinetic and safety data available in this age group. In children who need five drugs, delamanid is a preferred agent, although again, some clinical centers have reported good results with PAS. In children who are ages 3 to <6 years and who have fluoroquinolone resistant disease, delamanid is essential. If a fifth drug is needed, either ethionamide or PAS can be considered based on resistance pattern (i.e. ethionamide should not be given to persons with known or suspected inhA mutations) and tolerance.

Regimen design in children aged under 3 years can present some challenges and these recommendations are based mainly on clinical experience. Pharmacokinetic and safety data for many of the second-line drugs are limited in this population. Children in this age group with no evidence of fluoroquinolone resistance should be treated with Lzd-Lfx-Cfz-Cs (although there are limited data on the pharmacokinetics and safety of these drugs in this population, especially clofazimine and cycloserine). If a fifth drug is needed, clinicians will need to choose between delamanid, ethionamide and PAS, and there are no data to guide drug selection in this situation. Some providers will prefer to give delamanid since it has been assessed in pharmacokinetic and safety studies, and has a stronger body of evidence supporting its efficacy in adults. However, in pharmacokinetic studies, the selected doses did not result in adequate blood levels. Some clinical centers prefer to give PAS and have reported excellent outcomes with the use of this medication, although dosing and safety of PAS in children in this age group has also not been well established. Finally, ethionamide could be considered if there is no evidence of an inhA mutation. For children <3 years with fluoroquinolone resistance, linezolid, clofazimine and cycloserine should be given as well as at least one of delamanid, PAS or ethionamide. Others of these drugs can be added to strengthen the regimen. Ethambutol and/or pyrazinamide could be considered if susceptibility to these drugs is confirmed.

Figure 4. Severity of disease in children with TB



Case Examples: Regimen Design

Bangladesh

Rami is a 3-year-old boy who presents to the clinic with swollen “lumps” in his neck

His mother reports he has been sick for several weeks. He is not playing normally and is not eating well. She also reports him having fevers and a cough. She tells you that his auntie—who shares a bed with him—has TB and is being treated through a special program for people with drug-resistant TB. Rami’s physical examination and chest X-ray are consistent with TB. His rapid HIV test is negative. He is unable to cough sputum, even with induction, and two gastric aspirates are performed and sent for smear and culture. There is no bronchoscopy available where Rami lives. He is started on a short course of amoxicillin and told to return in 2 weeks.

Upon return to clinic, Rami has clinically worsened. He has lost 1 kg and his mother reports he sleeps all day. Results from his sputum smears were negative, and cultures are pending. He is barely arousable. A lumbar puncture is performed which shows no red cells but 15 nucleated cells of which 80% are lymphocytes. No AFB are seen in the spinal fluid.

You received results from his auntie’s physicians, which show that she has resistance to isoniazid, rifampicin, ethambutol and streptomycin.

Rami is diagnosed with TB meningitis and disseminated disease. Given his critical condition, he needs to start treatment immediately. He was likely infected by his auntie with whom he shares a bed, and she has documented drug resistance.

Rami is given five new medications based on the drug-resistance pattern of his source patient, consisting of Lfx-Lzd-Dlm-Cs-Ethio. Because Rami has TB meningitis, linezolid is an important drug in his regimen. Ethionamide is preferred over clofazimine since it has better central nervous system penetration. He is also started on prednisone as corticosteroids are indicated in cases of TB meningitis.

M tuberculosis never grows from his gastric aspirates and CSF samples. Rami improves and is continued on his regimen for a total of 18 months. Rami is then declared cured after 18 months of total therapy, based on his clinical improvement, resolution of chest x-ray findings, resolution of lymphadenopathy, and good weight gain.

Haiti

Angelie is a 12-year-old girl who is referred to the central hospital having failed a first-line anti-TB regimen. Angelie took her first-line medications daily under directly observed therapy. She had a positive sputum smear result at diagnosis, converted her sputum smear at month two, but had another positive sputum smear at month six. Samples were not sent for culture.

She reports feeling ill with cough, shortness of breath, and fevers. She is also losing weight. She reports that during the first month of treatment she felt “a little better” but overall feels worse now than she has since her diagnosis. She mentions that her father died of TB and that he had been in prison at the time. She was very close to him and visited him monthly.

Her clinical exam and X-ray are consistent with TB, and she has right upper lobe infiltrative lesions. Her sputum is AFB smear positive. It is sent for culture and for GeneXpert® testing. The GeneXpert® result is positive for *M. tuberculosis* and has detected rifampicin resistance. Her HIV rapid test is negative.

Angelie returns to the clinic 5 days later and now reports difficulty in breathing and wheezing. Her culture is pending, but given her contact history, her clinical worsening, and her GeneXpert® result, she is started on treatment for MDR-TB. Her father never had drug- susceptibility testing done.

She is started on four drugs to which she has not been previously exposed. She is started on Bdq-Lfx-Lzd-Cfz. She is also started on a bronchodilator inhaler for her wheezing.

Four weeks later, her DST results are obtained, showing her TB isolate has resistance to isoniazid and rifampicin but sensitivity to all other first and second-line medications. She is continued on a regimen of Bdq-Lfx-Lzd-Cfz. Because she is on linezolid, she has monthly monitoring of her complete blood count, monthly vision exams and a monthly examination for peripheral neuropathy. At her fourth month check, she has developed anemia with a hemoglobin of 9.2 g/dL. Her linezolid is stopped and cycloserine is initiated. After 6 months of negative cultures, her bedaquiline is discontinued, and she completes 15 months of therapy with Lfx-Cfz-Cs. She remains smear and culture negative, has gained weight, her symptoms have resolved, and after 15 months of therapy she is declared cured.

South Africa

JR is a 7-year-old boy whose mother was diagnosed with extensively drug-resistant TB (XDR-TB). When a post-exposure protocol is implemented and a nurse visits JR's home to see if anyone is sick there, she finds that JR has been coughing sputum for a month and has lost 3 kg. He also has fever and night sweats and no longer wants to play football with his teammates at school. He had an HIV test done 2 months ago which was negative.

On examination he is sweaty, pale, and warm to the touch. His weight is 26 kg. His conjunctiva are pale, he has no jaundice, he has no lymphadenopathy, and his heart exam is normal. His lung exam reveals crackles and bronchial breath sounds at the right apex. His abdomen is normal as are his extremities. A chest radiograph shows a large right upper lobe cavitory lesion. His sputum is sent for Xpert MTB/RIF® and is positive for both *M. tuberculosis* and rifampicin resistance. Because he has a known contact with XDR-TB, he is empirically started on an XDR-TB regimen while awaiting second-line DST.

He has a baseline ECG which reveals a QTc interval of 411msec and his baseline complete blood count is within normal limits. His initial regimen is Bdq-Lzd-Cfz-Cs-Dlm. His delamanid dose is 50mg twice a day as is recommended based on pharmacokinetic and safety studies and WHO recommendations. His bedaquiline dose is 200mg daily for 14 days followed by 100mg three times a week based on pharmacokinetic and safety studies and WHO recommendations.

He does well, gains weight, and is back playing football one month after starting treatment. Because he is on linezolid, he has monthly monitoring of his complete blood count, his vision, and for signs or symptoms of peripheral neuropathy.

Dosing of Second-Line Drugs in children

Proper dosing of second-line TB drugs in children is key to ensure good outcomes and to prevent the development of additional resistance. There are now pharmacokinetic (i.e. the way the body metabolizes a drug) and dosing data from pediatric populations and the doses of the second-line drugs recommended here are based on that information. Of note, dosing recommendations for very young children can be somewhat complicated (i.e. < 3 years). This is due to the fact that there are limited pharmacokinetic data on most second-line drugs in children, and optimal doses are yet to be determined.

KEY POINTS:

- Therapy in the absence of bacteriologic confirmation is more likely needed in children and should include a strong regimen that can be scaled back based on DST or development of adverse effects.
- Treatment should be based on the DST pattern of any known close contacts if the child does not have a DST of his or her own;
- Almost all children should be treated with injectable-free regimens;
- The duration of therapy in children should depend upon the extent of disease: children with non-severe disease can be treated for 9 to 12 months while children with severe disease will require 12-18 months of therapy depending on their clinical evolution;
- Regimens should consist of 4 to 5 presumed effective drugs for the duration of therapy: using more drugs adds to the toxicity of the regimen and does not likely improve efficacy if Group A and B drugs and/or delamanid are used: some groups have had good experience using PAS in children;
- Regimen construction should prioritize the WHO Group A and B drugs as well as delamanid;
- Monitoring and management of adverse events is essential;
- Child-friendly formulations of the medications should be used whenever possible.
- Corticosteroids should be used in cases of meningitis, pericarditis and disseminated disease.
- Linezolid should be given to all children with TB meningitis since its use has been associated with improved treatment outcomes.

(Photo credit Marcela Tommasi)



Ongoing pharmacokinetic studies will provide better data on the optimal use of second-line TB drugs in children of various age groups. This should include the newer drugs, bedaquiline and delamanid, as well as the repurposed agents, linezolid and clofazimine. Ideally this will be done in all relevant age groups, including children <3 years of age.

Another key challenge in ensuring optimal treatment for children is the limited access to child-friendly formulations of drugs used to treat MDR-TB in children. Table 5 documents the formulations that became available in 2018 for the treatment of MDR-TB via quality-assured mechanisms from the Global Drug Facility. Currently, most programs split adult tablets, which can lead to inconsistent dosing and violates Good Clinical Practice and Good Pharmacy Practice but may be the only option available. When available, child-friendly formulations should be prioritized.

There are emerging data on some of the second-line drugs in children. This manual provides the best data available now, to guide the management of children who need to be treated now. Further updates will be provided as more data are analyzed from this large study. Programs should use this while recognizing that updated dosing recommendations will be available soon.

Table 5. Formulations available from the Global Drug Facility

Drug	Formulation	Approval	Supplier
Ethambutol	100mg scored, dispersible tablet	WHO Pre-Qualified (PQ)	Via Global Drug Facility (GDF)
Isoniazid	100mg scored, dispersible tablet	Submitted	Via GDF
Pyrazinamide	150mg scored, dispersible tablet	WHO PQ	Via GDF
Levofloxacin	100mg scored, dispersible tablet	WHOPQ	Via GDF
Moxifloxacin	100mg scored, dispersible tablet	Expert Review Panel (ERP)	Via GDF
Linezolid	150mg scored, dispersible tablet	ERP	Not yet available
Ethionamide	125mg scored, dispersible tablet	WHOPQ	Via GDF
Cycloserine	125mg minicapsule	WHOPQ	Via GDF

Weight-Based Dosing in Children

GROUP A DRUGS

The following dosing tables (6a-6n) are meant to assist in optimal treatment of children with MDR-TB. The dosing ranges are based on the latest available pharmacokinetic data. However, it is important to realize that dosing recommendations can change quickly as additional studies are completed, and these recommendations could change. The weight-based dosing coincides with what is recommended by the WHO in their 2018 MDR-TB guidance. Some of the weight bands, however are different and the WHO starts their weight-bands at 5kg. This is because when the dispersible tablets are used, more precise dosing can be achieved within more narrow weight bands.

Table 6a: Levofloxacin

Levofloxacin 100mg scored, dispersible tablets Recommended dosing: 15-20mg/kg/day Weight-based dosing			
Weight Band (kg)	Dose	Number of 100mg tablets	Number of 250 mg tablets
1kg	20mg	Mix 100mg tablet in 10ml of water and administer 2ml of mixture immediately	-
2kg	40mg	Mix 100mg tablet in 10ml of water and administer 4ml of mixture immediately	-
3kg	50mg	0.5	-
4-6kg	100mg	1	0.5
7-9kg	150 mg	1.5	0.5
10-12kg	200-250mg	2.0 to 2.5	1
13-15kg	300mg	3	1-1.5
16-18kg	300-350mg	3-3.5	1.5
19-20kg	400mg	4	1.5
21-23kg	400-450mg	4-4.5	2
24-25 kg	500mg	5	2
26-35kg	750mg	-	3

Table 6b: Moxifloxacin

Moxifloxacin Recommended dosing: 10-15mg/kg/day Weight-based dosing			
Weight Band (kg)	Dose	Number of 100mg tablets	Number of 400mg tablets (dissolve in 10ml water)
1kg	10mg	Mix 100mg tablet in 10ml of water and administer 1ml of mixture immediately	-
2kg	20mg	Mix 100mg tablet in 10ml of water and administer 2ml of mixture immediately	-
3kg	30mg	Mix 100mg tablet in 10ml of water and administer 3ml of mixture immediately	-
4-6kg	50-80mg	0.5-0.75	2ml
7-9kg	150mg	1.5	3ml
10-15 kg	200mg	2.0	4ml
16-19 kg	300mg	3	0.5-.75 of a 400mg tablet
20-25kg	400mg	4	1
26-35kg	400mg	-	1

Table 6c: Linezolid

Linezolid Dose: 10-12 mg/kg/day > 16kg; 15mg/kg once daily in children < 16 kg Weight-based dosing				
Weight Band (kg)	Dose	150mg tablets (not yet available)	600mg tablet	20mg/ml suspension
1kg	15mg once daily	Mix 150mg tablet in 15ml of water and administer 1.5ml of mixture immediately	-	-
2kg	30mg once daily	Mix 150mg tablet in 15ml of water and administer 3 ml of mixture immediately	-	
3kg	45mg once daily	Mix 150mg tablet in 15ml of water and administer 4.5ml of mixture immediately	-	
4kg	60mg once daily	Mix 150mg tablet in 15ml of water and administer 6ml of mixture immediately	-	
5kg	75mg once daily	0.5 of 150mg tablet	-	4ml

Weight Band (kg)	Dose	150mg tablets (not yet available)	600mg tablet	20mg/ml suspension
6kg	90mg once daily	Mix 150mg tablet in 15ml of water and administer 9 ml of mixture immediately	0.25	4ml
7-9kg	75-150mg once daily	0.5-1.0 tablet	0.25	6ml
10-15kg	150-225mg once daily	1-1.5 tablet	0.25	8ml
16-20kg	225-250mg once daily	1.5-2 tablet	0.5	11ml
21-25kg	300mg once daily if < 12 years of age	2	0.5	14ml
36-35kg	300mg once daily if < 12 years of age	-	0.5	-

Table 6d: Bedaquiline

Bedaquiline		
Dose: 6mg/kg/day for 14 days followed by 3-4mg/kg thrice weekly (dose extrapolated from adult dosing for those less than 16 kg)		
Weight-based dosing		
Weight Band (kg)	Dose	100mg tablet
1-10kg	Consult with specialist ¹	
10-15kg	Consult with specialist <i>Likely recommended dose is 100mg daily for 14 days followed by 50 mg thrice weekly</i>	1 tablet daily for 14 days followed by 0.5 tablet thrice weekly (i.e. M/W/F)
16-23kg	200mg daily for 14 days followed by 100mg thrice weekly (i.e. M/W/F)	2 tablets daily for 14 days followed by 1 tablet thrice weekly (i.e. M/W/F)
24-30kg	200mg daily for 14 days followed by 100mg thrice weekly (i.e. M/W/F)	2 tablets daily for 14 days followed by 1 tablet thrice weekly (i.e. M/W/F)
31-34kg	400mg daily for 14 days followed by 200mg thrice weekly (i.e. M/W/F)	4 tablets daily for 14 days followed by 2 tablet thrice weekly (i.e. M/W/F)
> 34 kg	400mg daily for 14 days followed by 200mg thrice weekly (i.e. M/W/F)	4 tablets daily for 14 days followed by 2 tablets thrice weekly (i.e. M/W/F)

¹tbsentinelproject@gmail.com

GROUP B DRUGS

Table 6e: Clofazimine

Clofazimine Dosing: 2-5mg/kg/day Weight-based dosing			
Weight Band (kg)	Dose	50mg gelcaps	100mg gelcaps
<5kg	15mg	Give 1 gelcap M/W/F	Consult a specialist
5-6kg	10-30mg	Give 1 gelcap on alternative days	1 gelcap M/W/F
7-9kg	15-30mg	Give 1 gelcap on alternative days	1 gelcap M/W/F
10-15kg	20-75 mg	Give 1 gelcap either daily or on alternative days	1 gelcap alternative days
16-23kg	32-115mg	Give 1 gelcap per day	1 gelcap on alternative days
24-35kg	100 mg	Give 2 gelcap daily	1 gelcap daily

Table 6f: Cycloserine

Cycloserine Recommended dose: 15-20mg/kg/day Weight-based dosing			
Weight Band (kg)	Dose	125mg minicapsule	250mg capsule
1kg	20mg	Mix 125mg capsule in 12ml of water and administer 1ml of mixture immediately	-
2kg	40mg	Mix 125mg capsule in 12ml of water and administer 2ml of mixture immediately	-
3-4kg	62.5mg	Mix 125mg capsule in 12ml of water and administer 5ml of mixture immediately	-
5-9kg	125 mg	1	-
10-15 kg	250mg	2	1
16-23kg	375mg	3	2
24-35kg	500mg	4	2

GROUP C DRUGS (in order of how they should be used)

Table 6g: Delamanid

Delamanid Recommended dosing 3-4mg/kg/day (dose extrapolated from adult dosing for those less than 10 kg)		
Weight Band (kg)	Dose	50mg tablet
1-6kg	Consult with specialist ²	
7-23 kg	25 mg twice daily	½ tablet twice daily
24-34kg	50 mg twice daily	1 tablet twice daily
> 34 kg	100mg twice daily	2 tablets twice daily

²tbsentinelproject@gmail.com

Note that the 50mg tablet of delamanid when it is crushed, manipulated, or mixed does not result in the same blood levels as the 25mg pediatric formulation. Until the 25mg pediatric formulation is available, the 50mg tablet should be used with caution. Split tablets should not be saved for later administration for time periods longer than 12 hours.

According to recent WHO guidelines (2018), delamanid is preferred to bedaquiline in children <6 years, but bedaquiline can be considered in settings in which there is limited access to delamanid, although there are no data to guide the dosing of bedaquiline in this population and it would be considered an “expanded indication” (sometimes referred to as “off-label use”).

Table 6h: Ethambutol

Ethambutol 100mg Recommended Dosing: 15-25mg/kg/day Weight-based dosing			
Weight Band (kg)	Dose	100mg tablets	400mg tablets
1kg	20mg	Mix 100mg tablet in 10ml of water and administer 2ml of mixture immediately	-
2kg	40mg	Mix 100mg tablet in 10ml of water and administer 4ml of mixture immediately	-
3kg	70mg	Mix 100mg tablet in 10ml of water and administer 7ml of mixture immediately	-
4-6kg	100mg	1	-
7-9kg	200mg	2	-
10-12kg	250mg	2.5	-
13-15kg	300mg	3	-
16-18kg	350 mg	3.5	-
19-20kg	400mg	4	1
21-23kg	450mg	4.5	1
24-31kg	500mg	5	1.5
31-35kg	800mg	-	2

Table 6i: Pyrazinamide

Pyrazinamide Recommended dose: 30-40mg/kg/day Weight-based dosing			
Weight Band (kg)	Dose	150mg dispersible tablets	500mg tablet
1kg	30mg	Mix 150mg tablet in 10ml of water and administer 2ml of mixture immediately	-
2kg	60mg	Mix 150mg tablet in 10ml of water and administer 4ml of mixture immediately	-
3kg	90mg	Mix 150mg tablet in 10ml of water and administer 6ml of mixture immediately	-
4-6kg	150mg	1	-
7-9kg	225mg	2	-
10-12kg	375mg	2.5	-
13-15kg	450mg	3	-
16-18kg	525mg	3.5	1
19-20kg	600mg	4	1.25
21-23kg	675mg	4.5	1.5
24-30kg	750mg	5	1.5-2
31-35kg	1250mg		2.5

Table 6j: ethionamide

Ethionamide Recommended dose: 15-20mg/kg/day Weight-based dosing			
Weight Band (kg)	Dose	125mg tablets	250mg tablets
1kg	20mg	Mix 125mg tablet in 12ml of water and administer 1ml of mixture immediately	-
2kg	40mg	Mix 150mg tablet in 10ml of water and administer 4ml of mixture immediately	-
3-4kg	62.5mg	0.5	-
5-6kg	125mg	1	0.5
7-9kg	187.5mg	1.5	0.5
10-13 kg	250mg	2	1
14-15kg	312.5mg	2.5	1
16-20kg	375mg	3	2
21-23kg	437.5mg	3.5	2
24-30kg	500mg	4	2
31-35kg	500mg	-	2

Table 6k: PAS

Para-aminosalicylic acid (PAS either acid or sodium salt) 200-300mg/kg divided into 2 daily doses Some clinical centers give PAS 200mg/kg as a single daily dose and this could be considered Should be used with dosing spoon for more accurate dosing	
Weight Band (kg)	Dose
1kg	150mg twice daily
2 kg	300mg twice daily
3-4kg	500mg twice daily
5-6kg	0.5-0.75 gm twice daily
7-9kg	0.75-1.0 gm twice daily
11-13kg	1 gm twice daily
14-15kg	2 gm twice daily
16-20kg	2.5 gm twice daily
21-23kg	3 gm twice daily
24-30kg	3.5 gm twice daily
31-35kg	4gm twice daily

PASER® (PAS Acid) is stable for up to 8 weeks at 40°C and 75% humidity, and therefore can be distributed to the patient on a monthly basis in most environments with no cold chain. If storage of longer than 8 weeks is needed, refrigeration below 15°C is required.

Table 6l: Meropenem

Meropenem/Amoxicillin-clavulanate		
Drug	Daily dose	Maximum daily dose
Amoxicillin-clavulanate*	40mg/kg given twice daily based on the amoxicillin component	4000mg amoxicillin and 500mg clavulanate
Meropenem	20-40mg/kg IV every 8 hours	6000mg

*Amoxicillin-clavulanate should only be given in combination with meropenem. It should be given 30 minutes prior to the IV infusion of meropenem or imipenem.

Table 6m: Amikacin

Amikacin		
Drug	Daily dose	Maximum daily dose
Amikacin	15-20mg/kg once daily	1000mg

Amikacin should only be used in settings where susceptibility has been confirmed and where monthly, formal monitoring of hearing can be done (i.e. otoacoustic emissions in children <5 years of age, pure tone audiometry in children ages 5 years and above).

Table 6n: Isoniazid

Isoniazid 100mg* Recommended dosing: 15-20 mg/kg Weight-based dosing			
Weight Band (kg)	Dose	100mg dispersible tablets	300mg tablet
1kg	15mg	Mix 100mg tablet in 10ml of water and administer 1.5ml of mixture immediately	-
2kg	30mg	Mix 100mg tablet in 10ml of water and administer 3ml of mixture immediately	-
3kg	50mg	0.5	-
4-6kg	100mg	1	-
7-9kg	150mg	1.5	-
10-15kg	200mg	2	-
16-18kg	250mg	2.5	-
19-20kg	300mg	3	1
21-23kg	350 mg	3.5	1
24-30kg	400mg	4	1.5
31-35kg	600mg	-	2

*The role of HD-INH in the treatment of MDR-TB is still unclear but the drug could be considered in children with inhA mutations if there are no other options to construct an adequate regimen.

Pyridoxine should always be given with high dose isoniazid in children (12.5 mg daily in <5 year olds and 25 mg daily in >4 y olds).

Table 7: Central Nervous System Penetration of Second-Line Medications

Medication	CNS Penetration
Amikacin	Poor penetration except in the presence of meningeal inflammation
Bedaquiline	No data available; studies ongoing
Clofazimine	Limited data available
Cycloserine	CSF levels similar to serum levels
Delamanid	Limited human data but good CSF penetration in mice: studies ongoing
Ethambutol	Poor penetration
Ethionamide (prothionamide)	CSF levels similar to serum levels, but higher end dosing (20mg/kg) recommended in children
Isoniazid	20% of serum concentrations except in the presence of meningeal inflammation
Levofloxacin	65% of serum concentrations
Linezolid	Animal studies show CSF levels at 30% of serum levels: widely used in humans with excellent results
Meropenem	Excellent
Moxifloxacin	Good penetration in animals
PAS	Poor penetration except in the presence of meningeal inflammation
Pyrazinamide	CSF levels similar to serum levels

3. MONITORING

Diagnosing children with MDR-TB and designing an appropriate treatment regimen can be major obstacles in the management of pediatric MDR-TB. Another challenge is maintaining the patient on therapy for 9-18 months and making sure that he or she is closely followed by physicians, nurses, health care workers, and caregivers. Children have been successfully treated for MDR-TB, but only with appropriate monitoring and follow-up. Monitoring is needed to evaluate therapeutic efficacy and to mitigate the development of adverse events. This section will discuss:

- Timing and types of monitoring
- Adverse events and management strategies
- Management of co-morbid conditions
- Adherence support
- Nutritional monitoring and support

Table 8: Treatment monitoring schedule

All children	Baseline	Month										Ongoing	
		1	2	3	4	5	6	9	12	15	18		
HIV status	•												
Toxicity (symptoms, signs)	•	•	•	•	•	•	•	•	•	•	•	•	•
Height and weight	•	•	•	•	•	•	•	•	•	•	•	•	•
Audiology ¹	•	•	•	•	•	•	•						
Visual acuity testing	•	•	•	•	•	•	•	•	•	•	•	•	•
Color vision testing ²	•	•	•	•	•	•	•	•	•	•	•	•	•
CXR ³	•			•			•						
TB culture and DST ⁴	•	•	•	•	•	•	•	•	•	•	•	•	•
TSH, T4 ⁵	•			•			•	•	•	•	•	•	•
Hematology (FBC, diff) ⁶	•	•	•	•	•	•	•	•	•	•	•	•	•
LFTs, including AST, ALT and total bilirubin	•			•			•	•	•	•	•	•	
ECG to assess QTc interval ⁷	•	•	•	•	•	•	•						
Psychosocial counseling and adherence support	•	•	•	•	•	•	•	•	•	•	•	•	•

HIV-infected children	Baseline	Month										Ongoing
		1	2	3	4	5	6	9	12	15	18	
Cholesterol ⁸	•						•				•	•
CD4 count and viral load	•						•				•	•

¹ Monthly while on injectable and at 6 months following termination of injectable

² If on ethambutol

³ If any pulmonary involvement

⁴ Monthly if old enough to expectorate; if unable to expectorate and initially smear or culture positive, monthly until culture-converted then every three months; if initially smear and culture negative, perform if clinically indicated. It is NOT recommended to repeat gastric aspirates monthly for monitoring purposes. DST should only be done on specimens that are positive after a negative culture has been documented.

⁵ If on ethionamide, prothionamide or PAS

⁶ If on linezolid or HIV-infected

⁷ For children on bedaquiline, clofazimine, moxifloxacin or delamanid

⁸ For patients on ART, depending on the regimen

Table 9: Suggested Monitoring by Second-Line Medications (in alphabetical order)

Medication	Suggested Monitoring Tests (frequency as per Table 8)	Comments
Amikacin	Electrolytes, hearing, renal function,	Formal hearing testing must be done, including pure tone audiometry and/or otoacoustic emissions. If hearing loss cannot be formally assessed, amikacin should not be used
Bedaquiline	Electrolytes, liver function, QTcF interval	
Clofazimine	Electrolytes, liver function, QTcF interval	Patients and families should be counseled about skin color changes, since it can be distressing and/or lead to inadvertent disclosure.
Cycloserine	Psychosocial counseling	
Delamanid	Electrolytes, liver function, QTcF interval	
Ethambutol	Color vision, visual acuity	
Ethionamide/ prothionamide	Liver function, TSH/T4	
Isoniazid	Liver function, peripheral neuropathy	Should be administered with vitamin B6
Levofloxacin	Electrolytes, QTcF interval	Less likely to cause QTcF prolongation than moxifloxacin
Linezolid	Color vision, Complete blood count, peripheral neuropathy, visual acuity	
Meropenem		

Medication	Suggested Monitoring Tests (frequency as per Table 8)	Comments
Moxifloxacin	Electrolytes, QTcF interval	More likely to cause QTcF prolongation than levofloxacin
PAS	Electrolytes, liver function, TSH/T4	
Pyrazinamide	Liver function	

Safety Monitoring Using the New and Repurposed Drugs

The newer drugs (bedaquiline, delamanid) and the group A and B repurposed drugs (linezolid, clofazimine) are recommended for children with DR-TB (depending on the age group and the extent of disease and resistance) and these drugs have been shown to be safe and effective in adults. Providers may not be familiar with optimal strategies for monitoring and management of adverse events of these medications in children. Key issues that should be considered in children on these medications include:

- QTcF prolongation has been reported with clofazimine, bedaquiline, and delamanid (and also with the fluoroquinolones moxifloxacin and levofloxacin). Children should have their QTc intervals assessed at baseline and corrective strategies (i.e. electrolyte replacement, nutritional supplementation) implemented if needed. The QTcF values are the same in children as in adults. For males, a QTcF < 450msec is considered normal and in females a QTcF < 470msec is considered normal. Management of QTcF prolongation in children should follow the same steps as in adults, with symptom assessment, repetition of the ECG, electrolyte replacement, nutritional assessment, and review of other medications and possible clinical conditions. If the patient has symptoms or the QTcF is above 500msec, then TB drugs need to be withheld then possibly reintroduced, depending on the clinical situation of the child. Of note, it may be a challenge to use adult leads/electrodes in children given the size of the child's chest wall. If possible, pediatric leads should be used.
- Bone marrow toxicity manifesting itself as anemia, thrombocytopenia, or leukopenia may be seen in children on linezolid, and this marrow toxicity can progress rapidly. A baseline complete blood count should be checked and repeated after 2 weeks on linezolid then monthly. Any symptoms indicative of anemia, thrombocytopenia, or leukopenia should prompt measurement of a complete blood count. Any decrease of 1 or more grades in platelets or leukocyte counts should prompt weekly complete blood counts and discontinuation (at least temporarily) of linezolid until other causes can be assessed. Linezolid could be re-started at a lower dose if it is a key drug in the regimen. Anemia is commonly present in children with DR-TB (and also in children with HIV and those with malnutrition). Any decrease of 1 grade should prompt more frequent monitoring. Any symptomatic anemia or Grade 3 or 4 anemia should lead to discontinuation of linezolid until other causes can be assessed. Linezolid could be re-started at a lower dose if it is a key drug in the regimen.

Although there are different methods for grading severity of hematologic abnormalities, we recommend the following (based on the CTCAE version 5.0 available from https://ctep.cancer.gov/protocoldevelopment/electronic_applications/docs/ctcae_v5_quick_reference_8.5x11.pdf) in Table 10 below.

Table 10: Severity grading criteria for hematologic abnormalities

Abnormality	Grade 1 (mild)	Grade 2 (moderate)	Grade 3 (severe)
Hemoglobin	<lower limits of normal to 10 g/dL	8.0-9.9 g/dL	<8.0 g/dL
Platelet count	< lower limits of normal to 75,000 cells/uL	50,000-74,000 cells/uL	25,000-49,000 cells/uL
White blood cell count	Lymphocyte: <lower limits of normal to 800/ mm ³	500-799/ mm ³	200-499/ mm ³
	Neutrophil: < lower limits of normal to 1500/ mm ³	1000-1499/ mm ³	500-999/ mm ³

- Peripheral neuropathy can be caused by linezolid. All children and their caregivers should be asked about any symptoms of neuropathy or if the child is having problems with tripping, stepping on things, etc. All children should have a peripheral nervous system exam—especially reflexes—checked and documented at each visit. If there are signs or symptoms of peripheral neuropathy, then linezolid should be discontinued while other causes of neuropathy are investigated. Linezolid could be re-started at a lower dose if it is a key drug in the regimen.
- Optic neuritis can also be caused by linezolid. The first sign of this is usually a loss in visual acuity. Children should have their vision tested at baseline—using age appropriate tools, including picture charts, object tracking and papillary responses—and monthly while on linezolid. Any changes in visual acuity should be further investigated and linezolid held. Corticosteroids should be administered if there is concern for optic neuropathy as these can be vision sparing. Linezolid should be discontinued and not reintroduced if no other causes of the vision changes can be found.
- Skin color changes/hyperpigmentation are usually seen with clofazimine. Though this will resolve several months after treatment discontinuation, patients and their families should be counseled about this. The skin changes can be distressing and can lead to inadvertent disclosure of TB status.

Table 11: Identification and Management of Adverse Events

Type of adverse event	Likely culprit drugs	Identification	Management
Hepatotoxicity	Pyrazinamide, rifampicin, ethionamide/ prothionamide, bedaquiline, PAS, clofazimine, delamanid	Tender liver, visible jaundice	<p>Stop all drugs if ALT/AST > 5 times the upper limit of normal;</p> <p>Wait for liver function to return to normal; Re-introduce drugs one-by-one sequentially, every 2 days with monitoring of liver function before introducing the next drug. The least hepatotoxic drugs should be added first: E-Cfz-Lzd, Mfx/Lfx, Dlm. Then introduce the more hepatotoxic one by one every three days: Bdq, Eto-H-Z while monitoring liver function tests after each one to identify the responsible drug</p>
Visual problems	Ethambutol, linezolid	Regular (i.e. baseline and monthly) testing with Snellen and Ishihara Chart (or age appropriate measure including papillary responses and “fixate and follow” response in children < 2 years of age and symbol charts in children ages 3-5 years))	Stop ethambutol or linezolid (and do not re-introduce), refer the patient to ophthalmologist for further evaluation and management, start prednisone (1mg/kg/day with planned taper) and substitute with alternative drug.
Anemia	Linezolid	Monthly monitoring of CBC is essential for children on linezolid	<p>If moderate to severe, stop linezolid until anemia resolves. Could re-start at a lower dose (i.e. 10mg/kg once a day)</p> <p>A shorter course of linezolid could be considered in young children</p>
Thrombocytopenia	Linezolid	Monthly monitoring of CBC is essential for children on linezolid	<p>If moderate to severe, stop linezolid until thrombocytopenia resolves. Could re-start at a lower dose (i.e. 10mg/kg once a day)</p> <p>A shorter course of linezolid could be considered in young children</p>

Type of adverse event	Likely culprit drugs	Identification	Management
Leukopenia	Linezolid	Monthly monitoring of CBC is essential for children on linezolid	<p>If moderate to severe, stop linezolid until leukopenia resolves. Could re-start at a lower dose (i.e. 10mg/kg once a day)</p> <p>A shorter course of linezolid could be considered in young children</p>
Hearing problems	Amikacin, streptomycin,	Identified through audiometry or problems in communication	<p>Stop s the injectable drug if hearing loss > 26 dB (Grade 1), substituting with an alternative drug such as delamanid. Injectable agents should not be used if hearing loss cannot be formally monitored by audiometry</p>
Thyroid dysfunction	Ethionamide/ prothionamide, PAS	Regular blood testing (TSH) , clinical hypothyroidism or goitre	<p>Consider thyroxine supplementation if (a) clinical hypothyroidism, or (b) raised TSH and decreased FT4;</p> <p>Children clear thyroxine faster than adults, so daily replacement doses may be higher. Children (4-15 years): 4 mcg/kg/day (maximum dose is 200 mcg). Infants (1-3 years): 10-15 mcg/kg/day (maximum dose is 200 mcg).</p> <p>Monitor TSH every month and increase the dose by 25 mcg until TSH normalizes (TSH < 5 mIU/L). If raised TSH and normal FT4 repeat test in 1 month. Thyroid dysfunction resolves upon discontinuation of the cause agent. Hormone replacement must continue at least 2 to 3 months after completed DR-TB treatment.</p>
Electrolyte disturbances (Hypokalemia)	Amikacin, streptomycin, kanamycin, capreomycin	Regular blood testing (potassium)	<p>If potassium is low, replace with oral potassium and consider replacing magnesium as well. If potassium < 2.5 m eq hospitalize and replace IV.</p>

Type of adverse event	Likely culprit drugs	Identification	Management
Renal impairment	Amikacin, streptomycin,	Regular blood testing, symptoms of high potassium	If creatinine rises or potassium is elevated, stop injectable, substitute with alternative drug, dose three times a week or reduce dose.
Severe rash (SJS)	Any drug, although some drugs are more likely to cause rash, such as PZA	Severe rash, peeling mucus membranes, child unwell	Stop all drugs; Wait until clinical condition has improved; Re-introduce drugs one-by-one sequentially, every 2 days, monitoring clinically.
Nausea and vomiting	Ethionamide/ prothionamide, PAS	Clinically	Consider separating the dosing of ethionamide/prothionamide and also of PAS from the other drugs by giving it in the evening; Consider reducing the dose of ethionamide/prothionamide and building the dose up to full dose over 2 weeks. With new onset nausea and vomiting, should also consider hepatotoxicity, hepatitis, pancreatitis, or increased intracranial pressure.
Peripheral neuropathy	Isoniazid, linezolid, less frequently Cs	Clinically	Give pyridoxine. If clinically evident neuropathy, stop linezolid or isoniazid and substitute with another effective agent (i.e. delamanid). Could consider re-introducing linezolid at a lower dose but must monitor more frequently and discontinue linezolid should any signs or symptoms progress or recur. Can be challenging to monitor in young children, thus a shorter course of linezolid could be considered in young children.
Diarrhea	PAS	Clinically	Consider other causes; Encourage hydration Reduce dose; Consider drug substitution Consider loperamide if no blood in the stool or no fever.

Type of adverse event	Likely culprit drugs	Identification	Management
Neuropsychiatric problems	Terizidone, cycloserine, isoniazid, levofloxacin, moxifloxacin,	Seizures, headache, behaviour changes, depression, sleep disturbances	Verify correct dosing; Stop likely culprit drug; If symptoms persist, reintroduce and stop next most likely drug; If symptoms severe or persistent, stop all likely drugs or reduce dose.
Joint problems	Pyrazinamide, levofloxacin, moxifloxacin	Clinically	Verify correct dosing; Consider reducing dose/stopping possible culprit drug. Give non-steroidal anti-inflammatory drugs (NSAIs) e.g Ibuprofen. If acute swelling, redness and warmth are present in a joint, consider aspiration for diagnosis of gout, infections, autoimmune diseases, TB arthritis etc.
Painful injection sites	Amikacin, streptomycin,	Clinically	Add local anesthetic to drug in equal volumes; Vary site of injection on a daily basis; Consider stopping injection and substituting with a different agent; If severe, consider splitting dose and giving half into two different sites.
QTc prolongation	Moxifloxacin, bedaquiline, clofazimine, delamanid	Monthly assessment by ECG Fainting, racing heart, and severe chest pain A QTc interval is considered prolonged if it is > 500msec or if it is > 50msec and the patient has symptoms	Repeat the ECG; Check electrolytes and replace if needed, repeat the test Review ancillary drugs to see if any prolong the QTc interval; if so, stop likely culprit drug; Check thyroid (if hypothyroidism treat accordingly); Discontinue moxifloxacin and reassess; if still prolonged, discontinue clofazimine or bedaquiline.
Skin pigmentation/discoloration	Clofazimine	Skin turns a darker brown or orange color while on the medication, and may also become very dry	Reassure patient that this will improve 2-3 months after treatment stops and skin will return to normal color; If dry skin use hydrating creams.



Co-Morbid Conditions


Children with MDR-TB often suffer from other conditions. These may be pre-existing or develop as a result of their MDR-TB or may be in conjunction with their MDR-TB. Common co-morbid conditions seen in children with MDR-TB include HIV, diabetes mellitus, orthopedic problems, and reactive airway disease. In each of these cases, children do better in terms of MDR-TB outcomes when their co-morbid conditions are also aggressively treated and controlled. Management of co-morbid conditions should follow these principles:

- Management should occur at the same time as treatment for MDR-TB; waiting for MDR-TB treatment to finish or move to a “continuation” phase puts the patient at risk for poor outcomes from both conditions.
- Management should be provided in an integrated setting making care easy for the patient and reducing the risk of MDR-TB in other clinical settings (i.e. diabetes clinic, asthma clinic).
- Care should be taken to avoid giving drugs with overlapping toxicities when possible.

A detailed algorithm on the management of children with co-morbid HIV is included in this field guide. For other co-morbid conditions, we recommend the following:

- Diabetes mellitus: Blood sugar results may fluctuate in the setting of acute MDR-TB, and thus, more frequent monitoring of blood sugars is necessary. Medications for diabetes may need to be adjusted in this initial phase. In addition, common drugs used to treat TB may exacerbate glucose control problems and could have overlapping toxicities with both the disease itself (e.g. peripheral neuropathy) and with diabetes treatment regimens (e.g. oral antihyperglycemics). Patients may need to adjust their insulin dosing for tighter control, especially in the early stages of treatment. In addition, patients should be provided with adequate calories to ensure healthy weight gain.
- Reactive airway disease: Active MDR-TB can exacerbate existing reactive airway disease or cause reactive airway disease. Bronchodilators should be used for both maintenance and rescue situations. Inhaled corticosteroids can be safely used in children with MDR-TB.
- Orthopedic problems: Children may develop TB of the spine or joints, requiring the use of braces or other support devices. Children may also need physical therapy as part of their recoveries. When possible, local materials should be used for devices, and simple physical therapy regimens (e.g. chest clapping) that can be done at home should be designed.
- All children with MDR-TB should have a full complement of immunizations. It is important to verify immunizations at each appointment.

Figure 5: Algorithm for Management of Children on Treatment for MDR- TB and HIV



HAART: highly active antiretroviral therapy; MDR-TB: multidrug-resistant tuberculosis; CNS: central nervous system, IRIS: Immune Reconstitution Inflammatory Syndrome

Nutritional Monitoring for Children with MDR-TB

One of the key indicators for clinical monitoring in children being treated for MDR-

TB is improvement in nutritional status. If a child does not have an improvement in nutritional status while being treated for MDR-TB, that child has little chance of having a successful treatment outcome.

There are several ways to monitor the nutritional status of a child undergoing treatment for MDR-TB. A baseline measure of weight, height, and mid-upper arm circumference (MUAC) should be made in all children with MDR-TB. The MUAC is an indicator of acute malnutrition; if such malnutrition is present, acute nutritional

interventions are needed, according to local standards. At monthly follow-up, height and weight should be assessed. The weight-for-age and weight-for-height should be plotted for all children between 0 and 5 years, and body mass index (BMI) should be plotted for all children between 5 and 19 years. While there are many methods that can be used for assessing nutritional status, what is important in following MDR-TB patients is that their percentiles improve over time and do not decrease at any point in time.

If a child does not demonstrate improved nutritional status, this is a sign that his or

her MDR-TB is not being appropriately treated (or there may be another underlying clinical condition that needs to be diagnosed and treated). Children with MDR-TB require a higher caloric intake than their well counterparts because of the active metabolism associated with MDR-TB. Failure to improve nutritional status is an early and clear indicator that the MDR-TB may not be under control.

In addition to active MDR-TB contributing to poor nutritional status, many children and their families with MDR-TB often live in poverty. Thus, families may not be able to meet their basic nutritional needs. Nutritional support for children and their families

should be considered a routine part of clinical care for persons with MDR-TB. Physicians and nurses often counsel children and their parents that “the child needs to eat better.” This advice, however, is often not followed because the family cannot access food.

Nutritional counseling can still be given in this setting, and some practical advice is offered below.

Some programs try to improve nutritional status by prescribing vitamins for children with MDR-TB. These vitamins can be important sources of needed micronutrients, and vitamin B6 must be given to all children receiving therapy for MDR-TB. However, too many vitamins can increase the pill burden of the child and may not be well absorbed. It is always preferable to give the child vitamins combined with calories in the form of food. Ready to use therapeutic foods can be an important form of nutritional supplementation in children with MDR-TB. Programs should make an effort should be made to provide families with a food basket enough to feed the child and siblings to avoid excessive splitting of portions received by the MDR-TB patient alone.

Practical Nutritional Advice for Children and Families with MDR-TB

Children with MDR-TB and their families are often told that the child needs to “eat better.” They are given little, if any, practical advice on how to do so, especially in settings where they are unable to afford foodstuffs. This section offers practical advice on nutritional counseling for children and their families with MDR-TB.

- Know the resources in the community that offer nutritional assistance. TB programs may offer assistance directly to patients and their families on MDR-TB treatment. In other cases, there are additional groups working to provide nutritional support, such as non-governmental organizations, faith-based organizations, and community groups. These groups may give monthly or quarterly food provisions to families. Additional groups—including popular kitchens, community food groups, and food pantries—can provide single or ongoing cooked meals.
- Know the resources of the patient and family. Instead of telling them to “eat better,” ask them about the food they eat. Start by asking “how many meals do you eat a day?” Then proceed to ask about the composition of meals, who eats first in the family, and if there are any
- foods they avoid. Specifically ask if the child is able to drink milk.
- Know the locally available staple foods and the general price ranges for these foods. This can be done by taking a small shopping trip in the neighborhood. Once food availability and costs are known, patients and families can be given practical nutritional advice. For example, instead of encouraging them to “eat more protein” or “eat more meat,” recommend eggs (which contain protein but are often not as expensive as meat) or different cuts of meat (e.g. the liver or heart) which may cost less than other cuts. If the child is able to drink powdered milk, then the recommended recipe for making the milk could be “doubled” to increase caloric intake (i.e. add twice as much powder to the same amount of water). Nuts, legumes, and oil are all high protein foods that may be more affordable, depending on the setting. The same applies to leafy and green vegetables. Discourage families from buying expensive vitamin supplements and encourage them to invest instead in calorie- rich foods.



(Photo credit Marcela Tommasi)

- Recommend to the family that the child eat several small meals during the day. It can be overwhelming to a sick child to sit down to a larger plate of food. Eating multiple small, high calorie meals may help the child gain weight. This can be especially helpful for children with nausea and vomiting, which is common in patients on second-line anti-tuberculous therapy.

Adherence

Adherence to MDR-TB therapy is one of the cornerstones of treatment success. Hospitalization is not necessary for most children with MDR-TB and may actually decrease rates of adherence. As with adults, all children should be given treatment under DOT for each dose. Clinic-based DOT may place undue burdens on patients and their families, and where feasible, community-based DOT should be considered (using trained health workers provide treatment to the child in the household or community). In some situations, caregiver-administered therapy could be considered as well as long as there is effective monitoring and support for the parent/caregiver. If community-based DOT is not feasible, patients should be given incentives (e.g. food baskets) and enablers (e.g. transportation vouchers) to assist in treatment adherence and monthly follow up visits. Provision of DOT should go beyond “supervised swallowing” and include ongoing treatment literacy, monitoring for adverse events, and providing psychosocial support to children and their families.

Pediatric MDR-TB patients face special challenges with adherence. Very young children may not be able or willing to swallow tablets. Adolescent patients may use non-adherence as a way of asserting their independence. It is important to recognize that adherence strategies will need to be tailored to the individual patient and may change over time, even for the same patient. Some common principles should be followed in improving adherence among pediatric MDR-TB patients. These include:

- Age-appropriate patient education for the child and the caregiver. This is an extremely important part of adherence. The level of information given, and the manner in which it is delivered will need to be tailored according to the age of the child and where they are in their treatment course.
- Avoid the use of physical restraints and nasogastric tubes when possible. Avoidance may not be possible in all settings. Where restraints or nasogastric tubes are required, a daily assessment of ongoing need should be made.
- Adherence should be approached as a relationship, and pediatric patients offered some control over the process whenever possible (i.e. holding the medication spoon or dispenser with the provider; deciding the order in which to take the medications).
- It may be convenient to dose all medications at the same time, but this may be overwhelming for children. Twice daily dosing can be considered for some medications (e.g Cs, Ethio). Even with once daily dosing, half of the pills could be given in the morning and half in the evening, provided patients are not dosed the same medication more than once in 24 hours.
- Drug substitution to improve adherence (i.e. the changing of one effective medication for another in a successful treatment regimen to assist with adverse event management) can be considered, provided the substitution does not compromise the integrity of the regimen.

- Pediatric adherence depends on the caregivers. They should be involved at all stages and help make decisions about improving adherence.
- Incentives should be provided to children on a daily or weekly basis, depending on age. This could be as simple as a positive mark on a wall chart, singing a favorite song, or eating a special food. For adolescents, mobile phone minutes/data have been shown to be a powerful incentive. Incentives should also be provided to caregivers.

It is important to remember that children are often far more adherent than providers imagine them to be. Non-adherence may also be a sign of psychological or emotional distress, and social support should be given to both the child and the caregiver.

Case Examples: Adherence

Zimbabwe

Liswelicha is an 11-year-old boy diagnosed with MDR-TB. He is started on an MDR-TB regimen based on his DST which consists of Cfz-E-Bdq-Lfx-Lzd. He starts his medications in the hospital and initially does well, but after several episodes of vomiting, he refuses to take the “brown one” (i.e. clofazimine) because “it smells bad, makes my stomach hurt, and other kids make fun of my skin since I started that one.” He tries to run away from the hospital and once is found outside on the road looking for a taxi. He is adamant that the problem is the clofazimine, and he starts having anticipatory vomiting of all of his medications. Attempts are made to restrain and force him to swallow the clofazimine, but he continues to vomit and struggles strongly. A nasogastric tube is placed, but he removes it within an hour, even with both hands restrained.

A nurse talks with Liswelicha, and he again states that it is the “brown pill” he has a problem with. He says he will take all of his other medications—“even a shot”—as long as he does not have to take the clofazimine. A review of his DST confirms sensitivity to cycloserine, and the hospital has an adequate supply. His clofazimine is stopped, cycloserine is started, and Liswelicha becomes adherent, and his behavioral problems resolve.

Liswelicha is discharged from the hospital and does well for 7 months, with 100% adherence documented. He travels to a local clinic daily to take his medication. The clinic calls in his tenth month of treatment and says he has not come in for the past week. A home visit is made, and his mother notes he has moved to a camp for boys looking for work in the city. Liswelicha is found on a visit to the camp and states that he no longer needs his medication because he feels well. He also notes that now that he is 12, he needs to work to earn money for his family. A meeting is held with

Liswelicha and his mother, and he agrees to return home and finish his treatment. The clinic agrees to offer his family a monthly food basket for every month that Liswelicha remains on therapy. He completes 18 months of treatment then leaves to migrate to another country for work

Georgia

Nino is a 2-year-old, 10kg girl undergoing treatment for MDR-TB. Her father had confirmed MDR-TB and died while on therapy. Nino is in the hospital for her treatment, and she is started on Lzd-Lfx-Cs-PAS, as her father had resistance to isoniazid, rifampicin and ethambutol. Nino is quite ill and has major problems with vomiting her medications. Because she is in the hospital, her nurses and caregivers are able to space out her medications throughout the day, as the volume of the medications alone may be making her vomit. She has also developed diarrhea and soils her clothes at least three times a day. She cries and refuses to leave her room. Her mother is distraught and threatens to take Nino home.

The physicians review her medications and believe PAS is responsible, so they stop the drug and start her on delamanid at a dose of 25mg twice daily. A charity group brought some dolls to the hospital and Nino is enchanted with a small baby doll. The nurses use the doll to soothe Nino when she cries, and she develops a game where she pretends to give the doll medicine when she has to take hers. Nino still cries, but her doll provides a way for her to get comfort and to play act some of her fears. She completes 9 months of therapy, and both she and her doll are cured.

KEY POINTS:

- Children at different ages will have different adherence needs, and adherence needs change over time.
- Age-appropriate partnerships with children and their caregivers are key to improve adherence.
- Some adverse effects are serious (i.e. thrombocytopenia, neuropathy), and if treatment modifications can be made without affecting regimen integrity, these should be considered.
- Family factors that affect pediatric adherence should be addressed.

(Photo credit Jennifer Furin)



Special populations

Neonates

Neonates are at high risk for both TB infection and disease, include MDR-TB, and there are little data or experience in managing MDR-TB in these vulnerable children. In general, if a woman has MDR-TB and is culture positive in the last month of pregnancy, it can be presumed that the neonate has been exposed to MDR-TB. The baby can become sick with the disease, acquired either congenitally in utero or postnatally via airborne spread. Although there are limited data on the management of MDR-TB during pregnancy, clinical experience suggests that women can be treated with most of the commonly used MDR-TB drugs, including the novel agents, delamanid and bedaquiline. However, the injectables should be avoided, especially during the first trimester due to their adverse effects on the developing fetal ear. Therapeutic decisions regarding MDR-TB treatment during pregnancy must weigh the risks of untreated MDR-TB — including death of the mother and transmission to the neonate — versus the risks of exposing the neonate to the toxicity of second-line drugs. In the large majority of situations, treatment of MDR-TB during pregnancy is favored. Pyridoxine (vitamin B6) should be given to all pregnant women being treated for MDR- TB.

In terms of the management of babies born to women with MDR-TB, they should be evaluated for signs or symptoms of active disease at birth and then on a regular basis. The TST and IGRAs are of very limited utility in neonates. Complicating things, neonates are more likely to develop disseminated disease and exhibit non-classical TB symptoms, including irritability, poor feeding, splenomegaly, and hepatomegaly. To confirm a diagnosis of MDR-TB in a neonate whose mother had active MDR-TB during pregnancy, the placenta should be examined for pathologic indications of TB. Additional diagnostic studies in babies suspected of MDR-TB disease should include an immediate gastric aspirate and then 3 additional gastric aspirates, a lumbar puncture, blood samples, skin swabs and other relevant swabs/samples for mycobacteriologic assessment, with chest radiography or other imaging as indicated. If disease is suspected, treatment should be started as soon as possible and should be based on the DST of the mother. Dosing recommendations are the same as for older children.

Children born to women undergoing treatment for MDR-TB should be breast-fed if the mother is able to do so. Small concentrations of second-line drugs are present in the breast milk, so the neonate should be given pyridoxine (vitamin B6). Women who are still culture positive should wear a surgical mask when breastfeeding, and the baby should not share a bed with the mother. There is usually no need, however, to separate the mother-child pair as part of routine MDR-TB care, although separation may be considered in some situations for as short a period as possible. Sick and exposed neonates should be closely followed, preferably by a neonatologist or clinician with experience on MDR-TB in newborns. Preventive therapy should be given based on the drug susceptibility pattern of the mother, with fluoroquinolones being the ideal option. If preventive therapy is given, BCG vaccination should be postponed until such treatment is complete since these medications also kill the BCG and render it ineffective.

Adolescents

Adolescents (defined by the WHO as persons between the ages of 10 and 19 years) are often a population that can be overlooked when discussing pediatric MDR-TB. The clinical presentation of TB in this group and their ability to provide sputum for diagnostics often leads to them being thought of as adult patients. Some studies have shown that adolescents have worse treatment outcomes when compared with adults, especially those adolescents co-infected with HIV; others have demonstrated equivalent or even better outcomes when adolescents are compared with adults. While issues around adherence require special attention in this population with changing emotional and social needs, data show that adolescents may also have an increased death rate compared with adults. These deaths appear to occur early in the course of therapy, suggesting a delay in diagnosis. For these reasons, it is important to consider the needs of this special population in diagnosing MDR-TB and supporting adherence. The use of adolescent support clubs, such as those that have been used to support adolescents with HIV, could be of great utility in this population.

4. MANAGEMENT OF PERSONS EXPOSED TO MDR-TB IN THE HOUSEHOLD


The WHO now suggests that one option for the management of MDR-TB household exposure is the provision of MDR-TB preventive therapy to the contact. While randomized studies of the optimal treatment of such individuals are planned or ongoing, observational data show a clear benefit when treatment of MDR-TB infection (i.e. preventive therapy) is provided. Given the high risks associated with developing MDR-TB, a risk-benefit assessment of provision of such therapy favors its use. A detailed discussion of this is beyond the scope of this Field Guide, but can be found in the Sentinel Project document on how to care for persons exposed to MDR-TB (<http://sentinel-project.org/2018/03/29/how-to-care-for-people-exposed-to-drug-resistant-tuberculosis-a-practical-guide/>).

Following MDR-TB exposure, all household contacts should be evaluated for evidence of TB disease. If TB disease is ruled out, it is our opinion that every household member should then be offered MDR-TB preventive therapy. In some centers, only children under 5 years or those living with HIV are offered preventive therapy. In others, only children under 5 years, those living with HIV and those with a positive test of infection (TST and IGRA) are offered treatment. Whether treated or not, one of the most important aspects of the management of MDR-TB contacts is close follow up (probably every 3 months at a minimum) for at least a year after exposure to detect cases of incident TB early so that they can be treated early when disease has not been allowed to develop in severity.

In most cases, MDR-TB preventive therapy will be a third-generation fluoroquinolone such as levofloxacin or moxifloxacin. In a meta-analysis, fluoroquinolone-based prophylaxis was found to reduce the development of TB transmission by 90%. This can be given alone or in combination with another drug (such as ethambutol, high dose isoniazid or ethionamide). In cases of fluoroquinolone resistant TB, high dose isoniazid (15-20mg/kg) could be considered if the source patient has an inhA mutation. If not, delamanid could be considered under operational research conditions. There is no evidence to support the use of delamanid for the treatment of MDR-TB infection, but the drug is being tested in a clinical trial on the treatment of MDR-TB infection.

Optimal duration of MDR-TB preventive therapy is unknown, but our experience suggests that 6 months is appropriate. See Appendix C for Contact Management Form.

Figure 6. Suggested Management Algorithm for Child Contacts of MDR-TB Patients*



¹Infectious is defined as smear or culture-positive pulmonary TB

²Cough, reduced playfulness, fever, lethargy, bone or joint abnormalities, faltering weight

³If diagnosed by GeneXpert, consider MDR until confirmation by line probe assay (LPA) or DST
 CXR: chest x-ray; IGRA: interferon gamma release assay; TST: tuberculin skin test

Case Example: Contact

South Africa

Lelethu is a 2½-year-old girl who has been referred to your clinic, as she is a known contact of someone with MDR-TB. The person with MDR-TB is her uncle who lives in the same house as her but sleeps in a separate room. He had been treated with first-line therapy for 5 months before being diagnosed with MDR-TB and has been coughing for months. He spends most days with Lelethu because both her parents work during the day, and he is her main caregiver. He has now been on MDR-TB treatment for 2 months and is feeling much better. He is still getting injections daily from the clinic.

Lelethu's mother does not know much about his TB, so you ring the laboratory to get his results. You see that he had 3+ sputum smear-positive microscopy, and his TB is resistant to isoniazid and rifampicin but susceptible to the fluoroquinolones, amikacin and ethionamide.

Lelethu is very well, and during the consultation she rushes around the clinic room playing with everything, and she appears very happy. Her mother tells you that she is not coughing and has no fever or sweating. You measure her weight, height and mid-upper arm circumference (MUAC) and then ask to look at her road-to-health card.

She seems to be growing very well along the 25th percentile. Clinical examination is completely normal.

You decide that she should receive MDR-TB preventive treatment with isoniazid at high dose (15-20mg/kg) since her contacts have inhA mutations (meaning high-doses of INH should be effective) and levofloxacin (15-20mg) daily for 6 months. You also request an HIV test (even though her mother says that she was tested in pregnancy and was found to be negative). The HIV test is negative. Lelethu is reviewed by the nurse every month to measure her weight, check how she is getting on with the medications, and to ask if there are any problems. Every 3 months Lelethu comes back to your clinic to be seen by a nurse and a physician. She takes her preventive therapy for 6 months, and at the twelfth month, she is still fit and well. Her weight and height have increased. She is then discharged.

5. INFECTION CONTROL

Although most children with TB are not contagious, infection control is of paramount importance in the management of MDR-TB in children. Children should be protected from becoming infected with MDR-TB in both the health facility and home setting. Children with MDR-TB should be safely managed in a way that does not cause unnecessary psychosocial stress and avoids making them victims of stigma. Children with MDR-TB usually do better in a home setting and when they are able to resume normal activities, such as going to school. In most cases, as long as the child is on appropriate therapy for MDR-TB, the risk of transmitting MDR-TB is low. This section offers practical guidance on facility-based infection control and home/community-based infection control that acknowledges the need to reduce MDR-TB transmission risk while at the same time acknowledging the important developmental needs of a growing child. This section will discuss:

- Screening of household members
- Facility-based infection control
- Community-based infection control

Screening of household members

It is an urgent priority for TB programs to provide screening of any household member exposed to MDR-TB. For details on how to implement effective screening programs, please refer to the Sentinel Project's handbook on management of persons exposed to MDR-TB in the household (<http://sentinel-project.org/2018/03/29/how-to-care-for-people-exposed-to-drug-resistant-tuberculosis-a-practical-guide/>). Of note, if a child is diagnosed with MDR-TB, all household members should be screened to try and identify a possible source patient who can also be started on treatment.

Facility-Based Infection Control

Although negative pressure airflow isolation rooms and precautions are the gold standard in TB infection control, there are simple infection control measures that can be easily put into place to make nosocomial transmission of MDR-TB less likely. These include:

- Having patients wait outdoors;
- Using windows for natural ventilation;
- Having separate waiting areas for TB and MDR-TB patients with separate entrances and air supplies;
- Considering separate waiting areas for patients with cough if space allows;
- Separating waiting areas for ART clinics and TB clinics;
- Ensuring airflow is away from the health care provider during the consultation;
- Re-arranging consultation area furniture to follow the direction of airflow (i.e. to ensure health care providers and general waiting areas are away from the direction of contaminated airflow);
- Avoiding scheduling patients for well visits on days when known MDR-TB patients are being seen;
- Ensuring that appropriate therapy be given and maintained for all TB patients;
- Having patients with active cough wear surgical masks to decrease transmission;
- Avoiding unnecessary hospitalizations;
- Discharging patients on treatment from the wards as quickly as possible once effective therapy has been started and can be maintained in the community. This can be within days to weeks of starting MDR-TB therapy;
- Taking special infection control measures during highly infectious diagnostic procedures such as induced sputum collection;
- Consider other environmental infection control measures such as the use of ultraviolet lights.

Case Example: Infection Control

Rwanda

At a primary care clinic in rural Rwanda, the nurse notes that three patients have been diagnosed and are now being treated for MDR-TB. One of them is a 5-year-old child who was seen at the clinic 6 months ago with diarrhea. The nurse is very worried about TB transmission in her small, four-room clinic. She requests some benches from the government so that patients can wait outside, but she knows this will not work in the rainy season. She runs an antenatal clinic on Wednesday mornings and a vaccination clinic on both Wednesday and Thursday mornings. She asks all TB patients to come for sputum testing and DOT after 2 p.m. on Wednesdays and Thursdays, and she leaves

the clinic windows open at night. She also arranges for home-based DOT for her MDR-TB patients and visits them in their homes monthly to perform follow-up visits and collect specimens. She also requests a small awning for the clinic, so that all sputum specimens can be collected outside, even in the rain.

Community-Based Infection Control

Patients with MDR-TB can be safely treated in the community setting, and the risk for ongoing transmission is low, once the patient is on appropriate MDR-TB therapy. Some community and household measures should be taken to decrease transmission in the household and community. These include:

- MDR-TB patients should ideally sleep in a separate room.
- Windows in the home of a MDR-TB patient should be kept open as often as possible.
- MDR-TB patients should spend as much time outside whenever possible, including visits with friends and family members.
- MDR-TB patients should be provided with social support to be able to stay on MDR-TB therapy.

Household and community members often fear becoming infected with MDR-TB when a child with MDR-TB is returned to the community. As long as the child is maintained on appropriate treatment for MDR-TB, his or her risk of infectiousness is low. Once his or her smear is negative, he or she should return to normal activities—including school and sporting teams—provided his or her clinical status allows. Education should be provided to family members and key community members (i.e. teachers, coaches, ministers). This will decrease stigma and discrimination. Specific points to address include:

- Once the child is on MDR-TB treatment, he or she is unlikely to transmit disease.
- If smear-negative, the child doesn't need to wear a mask in public.
- Children with MDR-TB can share bathrooms, utensils, balls, tools, crayons, etc.
- Children with MDR-TB feel better physically and psychologically when they can return to their usual environments and activities.

Case Example: Infection Control

Peru

Blanca is a 9-year-old girl who was diagnosed with MDR-TB 3 months ago. She is clinically improved and ready to return home. Her family and friends are glad she is coming back, but they are worried they might get sick from her. They begin to gossip in the neighborhood, and some people even threaten to block her from coming into her apartment building.

Worried, her mother speaks to the local priest who agrees to help her talk about MDR-TB with the community. He preaches a sermon the Sunday before Blanca comes home in which he gives the community facts about MDR-TB. He and Blanca's mother answer any questions people have. Blanca's apartment is small, and she cannot have her own sleeping area. However, the church donates a mattress to her mother, and Blanca returns home to sleep there. She returns to school a week after coming home. Her teacher sends a note home her second week back stating that Blanca was punished for borrowing another child's pencil. Blanca's mother arranges for the priest to speak with the school as well. Blanca receives her DOT from a community health worker prior to going to school every day. After 2 months, she is thriving in the community and playing on the girls' football team.

Photo credit: Marcela Tommasi



SELECTED REFERENCES

This field guide is meant to provide practical advice for clinicians and programs managing children exposed to, infected with and sick with MDR forms of TB. Much of the material in this handbook comes from collected expert opinion. In addition to expert opinion, some key references were used in the development of this guide. These include:

Peer-Reviewed Articles

Al-Dabbagh M, Lapphra K, McGloin R, Inrig K, Schaaf HS, Marais BJ, et al. Drug-resistant tuberculosis: pediatric guidelines. *Pediatr Infect Dis J*. 2011 Jun;30(6):501-5.

Cohen T, Murray M, Abubakar I, Zhang Z, Sloutsky A, Arteaga F, et al. Multiple introductions of multidrug-resistant tuberculosis into households, Lima, Peru. *Emerging Infectious Diseases*. 2011 Jun;17(6):969-75.

Cruz AT, Garcia-Prats AJ, Furin J, Seddon JA. Treatment of Multidrug-Resistant Tuberculosis Infection in Children. *Pediatr Infect Dis J*. 2018 Oct; 37(10):1061-4.

Dheda K, Chang KC, Guglielmetti L, Furin J, Schaaf HS, Chesov D, et al. Clinical management of adults and children with multidrug-resistant and extensively drug-resistant tuberculosis. *Clin Microbiol Infect*. 2017;23(3):131-40

Ettehad D, Schaaf HS, Seddon JA, Cooke GS, Ford N. Treatment outcomes for children with multidrug-resistant tuberculosis: a systematic review and meta-analysis. *Lancet Infect Dis*. 2012 Jun;12(6):449-56.

Golla V, Snow K, Mandalakas AM, Schaaf HS, Du Preez K, Hesselning AC, et al. The impact of drug resistance on the risk of tuberculosis infection and disease in child household contacts: a cross sectional study. *BMC infectious diseases*. 2017;17(1):593.

Haraus EP, Garcia-Prats AJ, Law S, Schaaf HS, Kredo T, Seddon JA, et al. (2018) Treatment and outcomes in children with multidrug-resistant tuberculosis: A systematic review and individual patient data meta-analysis. *PLoS Med* 15(7): e1002591. <https://doi.org/10.1371/journal.pmed.1002591>

Haraus EP, Garcia-Prats AJ, Seddon JA, Schaaf HS, Hesselning AC, Achar J, et al. New and Repurposed Drugs for Pediatric Multidrug-Resistant Tuberculosis. Practice-based Recommendations. *Am J Respir Crit Care Med*. 2017;195(10):1300-10.

Jenkins HE, Tolman AW, Yuen CM, et al. Incidence of multidrug-resistant tuberculosis disease in children: systematic review and global estimates. *Lancet*. 2014;383:1572–9

Jonckheree S, Furin J. Overcoming challenges in the diagnosis, prevention, and treatment of pediatric drug-resistant tuberculosis. *Expert Rev Respir Med*. 2017;11(5):385-94.

Marks SM, Mase SR, Morris SB. Systematic Review, Meta-analysis, and Cost-effectiveness of Treatment of Latent Tuberculosis to Reduce Progression to Multidrug-Resistant Tuberculosis. *Clin Infect Dis*. 2017 Jun 15;64(12):1670-1677. doi: 10.1093/cid/cix208.

McAnaw SE, Hesselning AC, Seddon JA, Dooley KE, Garcia-Prats AJ, Kim S, et al. Pediatric multidrug-resistant tuberculosis clinical trials: challenges and opportunities. *Int J Infect Dis.* 2017;56:194-9.

Nachman S., Ahmed A., Amanullah F., et al. Towards early inclusion of children in tuberculosis drugs trials: a consensus statement. *Lancet Infect Dis.* 2015 Jun; 15(6):711-20

Reubenson G. Pediatric drug-resistant tuberculosis: a global perspective. *Paediatr Drugs.* 2011 Dec 1;13(6):349-55.

Schaaf HS, Marais BJ. Management of multidrug-resistant tuberculosis in children: a survival guide for paediatricians. *Paediatr Respir Rev.* 2011 Mar;12(1):31-8.

Schaaf, HS., Garcia-Prats, A., McKenna, L. and Seddon, J. Challenges of using new and repurposed drugs for the treatment of multidrug-resistant tuberculosis in children. *Expert Review in Clinical Pharmacology* 2018, 11(3): 233-44.

Seddon JA, Godfrey-Faussett P, Hesselning AC, Gie RP, Beyers N, Schaaf HS. Management of children exposed to multidrug-resistant *Mycobacterium tuberculosis*. *Lancet Infect Dis.* 2012 Jun;12(6):469-79.

Seddon JA, Hesselning AC, Finlayson H, Fielding K, Cox H, Hughes, et al. Preventive therapy for child contacts of multidrug-resistant tuberculosis: a prospective cohort study. *Clin Infect Dis.* 2013 Dec;57(12):1676-84.

Walters, E., van der Zalm, M., Palmer, M., et al. Xpert MTB/RIF on stool is useful for the rapid diagnosis of tuberculosis in young children with severe pulmonary disease. *Pediatric Infectious Diseases Journal* 2017, 36(9): 837-43

Books, Conference Proceedings, and Clinical Guidelines

Curry International Tuberculosis Center, California Department of Public Health. Drug-resistant tuberculosis: A survival guide for clinicians, 2011. [Accessed at: http://www.nationaltbcenter.ucsf.edu/drtb/docs/MDRTB_book_2011.pdf]

European Center for Disease Prevention and Control. Management of contacts of MDR TB and XDR TB patients. Stockholm: European Center for Disease Prevention and Control; 2012. [Accessed at: <http://ecdc.europa.eu/en/publications/Publications/201203-Guidance-MDR-TB-contacts.pdf>]

Institute of Medicine (US), Academy of Science of South Africa. Drug-resistant TB in children. The emerging threat of drug-resistant tuberculosis in Southern

Africa: Global and local challenges and solutions: Summary of a joint workshop, Washington, D.C.: National Academies Press (US); 2011. [Accessed at: <http://www.ncbi.nlm.nih.gov/books/NBK55582/pdf/TOC.pdf>]

Médecins Sans Frontières. Treating drug-resistant tuberculosis: what does it take? Médecins Sans Frontières; 2011. [Accessed at: http://www.msf.org.uk/TB_Report_2011.aspx]

Partners In Health. PIH guide to the medical management of multidrug-resistant tuberculosis. Boston: Partners In Health; 2003. [Accessed at: [http:// parthealth.3cdn.net/9512565bbaca05aca2_aqm6v2gyk.pdf](http://parthealth.3cdn.net/9512565bbaca05aca2_aqm6v2gyk.pdf)]

Sentinel Project on Pediatric Drug-Resistant Tuberculosis. Rapid clinical advice: The use of delamanid and bedaquiline for children with drug-resistant tuberculosis. Boston: Sentinel Project on Drug-Resistant Tuberculosis. 2016. [Accessed at:

[http://sentinel-project.org/2016/05/16/advancing-access-for-new-tb-drugs-for- children/](http://sentinel-project.org/2016/05/16/advancing-access-for-new-tb-drugs-for-children/)]

Treatment Action Group. HIV, TB, HCV Pipeline Report, 2018. Available at www.treatmentactiongroup.org/pipeline-report.

World Health Organization. WHO treatment guidelines for multidrug- and rifampicin-resistant tuberculosis: 2018 Update. Pre-Final Text. Available from: <https://www.who.int/tb/publications/2018/WHO.2018.MDR-TB.Rx.Guidelines.prefinal.text.pdf>.

World Health Organization. Rapid Communication. Key changes to the treatment of multidrug- and rifampicin-resistant tuberculosis. August 2018. https://www.who.int/tb/publications/2018/WHO_RapidCommunicationMDRTB.pdf.

World Health Organization. WHO treatment guidelines for multidrug- and rifampin-resistant tuberculosis. 2018 update. Available at: <https://www.who.int/tb/publications/2018/WHO.2018.MDR-TB.Rx.Guidelines.prefinal.text.pdf>.

World Health Organization. Guidelines for the programmatic management of drug-resistant tuberculosis: Emergency update 2008. Geneva: World Health Organization; 2008. [Accessed at: [http://whqlibdoc.who.int/ publications/2008/9789241547581_eng.pdf](http://whqlibdoc.who.int/publications/2008/9789241547581_eng.pdf)]

World Health Organization. Management of MDR-TB: A field guide. Geneva: World Health Organization; 2008. [Accessed at: [http://whqlibdoc.who.int/ publications/2009/9789241547765_eng.pdf](http://whqlibdoc.who.int/publications/2009/9789241547765_eng.pdf)]

World Health Organization, 2014. Guidance for national tuberculosis programmes on the management of tuberculosis in children. http://apps.who.int/iris/bitstream/handle/10665/112360/9789241548748_eng.pdf?sequence=1

World Health Organization. Towards universal access to diagnosis and treatment of multidrug-resistant and extensively drug-resistant tuberculosis by 2015 Geneva, Switzerland. ISBN: 978 92 4 150133 . WHO/HTM/TB/2011.3. 2011.

APPENDIX A: SAMPLE INTAKE FORM

PERSONAL DATA	
Name:	Date of evaluation:
Caregiver:	Date of birth:
Place of Residence:	Age:
	Sex:
	Medical record number:
Telephone:	
Address and mobile phone number (if different than above):	
Name of evaluator:	
Health establishment:	
TB HISTORY	
<input type="checkbox"/> Never diagnosed	Year first diagnosed with TB:
Ever received BCG?	Diagnosed by:
<input type="checkbox"/> Yes, year(s):	<input type="checkbox"/> AFB <input type="checkbox"/> Other (specify):
<input type="checkbox"/> No	<input type="checkbox"/> Culture
<input type="checkbox"/> Unknown	<input type="checkbox"/> CXR
Suspect primary MDR-TB	
<input type="checkbox"/> Yes <input type="checkbox"/> No	
Check all risk factors that apply:	
<input type="checkbox"/> Close contact with known MDR-TB	
<input type="checkbox"/> Close contact with person who died of TB or failed TB treatment	
<input type="checkbox"/> Previous treatment	
<input type="checkbox"/> Failure to improve on current TB treatment	
Summary of previous antituberculous drug use (mark each drug patient has received for >1 month)	
<input type="checkbox"/> INH	<input type="checkbox"/> SM
<input type="checkbox"/> RIF	<input type="checkbox"/> KM
<input type="checkbox"/> EMB	<input type="checkbox"/> AMK
<input type="checkbox"/> PZA	<input type="checkbox"/> CM
<input type="checkbox"/> FQ	<input type="checkbox"/> AMX-CLV
<input type="checkbox"/> THA/PTO	<input type="checkbox"/> CFZ
<input type="checkbox"/> CS	<input type="checkbox"/> Other:
<input type="checkbox"/> PAS	<input type="checkbox"/> Other:
IMMUNIZATION HISTORY	
Has the child been fully immunized for age:	Has patient had BCG?
<input type="checkbox"/> Yes	<input type="checkbox"/> Yes
<input type="checkbox"/> No	<input type="checkbox"/> No
If no, what vaccines are missing?	Is BCG scar present?
	<input type="checkbox"/> Yes
	<input type="checkbox"/> No

SOCIO-DEMOGRAPHIC DATA

Currently in school: <input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> N/A	Who are the primary caregivers? Are the caregivers employed? <input type="checkbox"/> Yes <input type="checkbox"/> No If yes, what is their work?
Number of household members:	
Number of household members when diagnosed with TB:	
Number of household members when diagnosed with MDR-TB:	
How far does patient live from health facility?	Have parent(s) been tested for HIV? <input type="checkbox"/> Yes <input type="checkbox"/> No If yes, date and result:
How did patient get to the health facility?	
How long does it take patient to get to the health facility?	

REVIEW OF SYSTEMS

Check all that apply	
<input type="checkbox"/> Cough <input type="checkbox"/> Sputum <input type="checkbox"/> Poor appetite <input type="checkbox"/> Weight loss <input type="checkbox"/> Bronchospasm	<input type="checkbox"/> Dyspnea at rest <input type="checkbox"/> Dyspnea on exertion <input type="checkbox"/> Tired, listless, not playing
<input type="checkbox"/> Swelling or "lumps" in neck, arms or groin <input type="checkbox"/> Vertebral pain <input type="checkbox"/> Back Pain <input type="checkbox"/> Other:	
<input type="checkbox"/> Hemoptysis Largest quantity in mL:	Date of first episode of hemoptysis:
Most recent quantity in mL:	Date of most recent hemoptysis:
Current medications:	Allergies or adverse reactions:

PAST MEDICAL HISTORY			
Diabetes <input type="checkbox"/> Yes <input type="checkbox"/> No	Asthma <input type="checkbox"/> Yes <input type="checkbox"/> No	Previous hospitalization(s)? <input type="checkbox"/> Yes <input type="checkbox"/> No Hospitalization(s) in pulmonary ward? <input type="checkbox"/> Yes <input type="checkbox"/> No Reason for hospitalization(s): Name of hospital(s):	
Other:			
Has patient been tested for HIV? <input type="checkbox"/> Yes <input type="checkbox"/> No If yes, date and result:			
Prior transfusion(s) <input type="checkbox"/> Yes <input type="checkbox"/> No		Date of transfusion(s): Indication for transfusion(s):	
BIRTH HISTORY AND PAST SURGICAL HISTORY			
Was patient born at home? <input type="checkbox"/> Yes <input type="checkbox"/> No Did the mother receive prenatal care? <input type="checkbox"/> Yes <input type="checkbox"/> No Were there any problems at birth? <input type="checkbox"/> Yes <input type="checkbox"/> No If yes, describe:		Prior surgery? <input type="checkbox"/> Yes <input type="checkbox"/> No Procedure(s): Date(s) of surgery: Complications:	
PHYSICAL EXAM			
Temp:	BP:	HR:	RR:
Weight:	Height:	BMI:	

GENERAL APPEARANCE

<p>HEENT</p> <p>Conjunctiva:</p> <p>Sclera:</p> <p>Oropharynx:</p>	<p>Lymphadenopathy present?</p> <p>JVD:</p> <p>Thyromegaly:</p>	<p>Cor</p> <p>Tachycardic?</p> <p>Murmurs?</p> <p>Extra heart sounds?</p>	<p>Lungs</p> <p>Wheezing?</p> <p>Crackles?</p> <p>Bronchial breathing sounds?</p>
<p>Abdomen</p> <p>Bowel sounds?</p> <p>Organomegaly?</p> <p>Tender?</p>	<p>Extremities</p> <p>Edema?</p> <p>Cyanosis?</p> <p>Pulses:</p>	<p>Neuro</p> <p>Mental status:</p> <p>Reflexes:</p> <p>Strength:</p> <p>Gait:</p>	<p>Developmental</p> <p>Describe development for age:</p>

TEST RESULTS

Drug susceptibility testing:

	Sample number	Date of sample collection	Date of results	AFB results (Pos, Neg, Unknown)	Laboratory	Resistant to	Susceptible to
1				P N U			
2				P N U			
3				P N U			
4				P N U			
5				P N U			
6				P N U			

Chest radiograph:

Other lab results:

Impression/plan:

SUMMARY OF KNOWN TB CONTACTS



Name of contact	Relation to patient	Date of TB diagnosis	Lived in same household when contact had TB?	History of multiple treatments?	Died during treatment?	History of documented MDR-TB?	Current AFB status*	Current status of TB contact**	Resistant to which drugs?
			Y N U	Y N U	Y N U	Y N U	P N U	C T S D U	
			Y N U	Y N U	Y N U	Y N U	P N U	C T S D U	
			Y N U	Y N U	Y N U	Y N U	P N U	C T S D U	
			Y N U	Y N U	Y N U	Y N U	P N U	C T S D U	

Unless stated otherwise, indicate **Y**es, **N**o or **U**nknown

For AFB status, indicate **P**ositive, **N**egative or **U**nknown

** For current status, indicate **C**ured, in **T**reatment, **S**ymptomatic but not in treatment, **D**eceased, or **U**nknown

APPENDIX B: MEDICATIONS USED TO TREAT MDR-TB

<p>Bedaquiline (BDQ) Tablet, 100mg</p>	
<p>Delamanid (DLM) Tablet, 50mg</p>	
<p>Linezolid (LZD) Tablet, 600mg</p>	
<p>Levofloxacin (LFX) 250mg tablet, 100mg dispersible tablet</p>	
<p>Ethionamide (THA) Form: tablet Dose: 250 mg or 125mg dispersible tablet</p>	
<p>Amoxicillin-clavulanic acid (AMX-CLV) Form: tablet Dose: 500 mg</p>	

<p>Moxifloxacin (MFX) Form: tablet Dose: 400 mg or 100 mg dispersible tablet</p>	
<p>Clofazimine (CFZ) Form: soft gel Dose: 100 mg, 50mg</p>	
<p>Cycloserine (CS) Form: capsule Dose: 250 mg or 125mg capsule</p>	
<p>Amikacin 500mg vial, IM</p>	
<p>Para-Aminosalicylic acid (PAS) Form: granules Dose: 4 g</p>	

APPENDIX C: CONTACT MANAGEMENT FORM FOR CHILDREN EXPOSED TO MDR-TB

CHILD PERSONAL INFORMATION			
Child name:		Clinic name:	
Child date of birth:		Clinic phone number:	
Child folder number:		Clinic fax number:	
Child address:			
SOURCE PATIENT INFORMATION			
Source patient name:		Date of sample production:	
Relationship to child:		Sputum smear result:	
DST results:			
	Resistant	Susceptible	Not tested
Isoniazid			
Rifampicin			
Fluoroquinolones			
Amikacin			
Ethionamide			
PHYSICAL EXAM			
Weight:	HIV test date:	Mantoux test date:	CXR date:
Height/length:	HIV test result:	Mantoux test size (mm):	CXR impression:
Symptoms:		Management:	
<input type="checkbox"/> Cough > 2 weeks <input type="checkbox"/> Losing weight <input type="checkbox"/> Night sweating	<input type="checkbox"/> Fever <input type="checkbox"/> Reduced energy <input type="checkbox"/> Abnormal joints/spine	<input type="checkbox"/> Refer <input type="checkbox"/> Prophylaxis <input type="checkbox"/> Discharge/observe	

PREVENTIVE TREATMENT

	Date started	Dose	Number of tablets
Isoniazid			
Levofloxacin			
Ethambutol			
Other:			

MONITORING CHART

Month	Weight	Height	Clinical review completed
1			
2			
3			
4			
6			
9			
12			

APPENDIX D: PROTOCOLS FOR SPECIMEN COLLECTION

Gastric Aspiration

Gastric aspiration can be used in children when sputa cannot be spontaneously expectorated nor induced using hypertonic saline. Since gastric aspiration is not an aerosol-generating procedure, it poses a low risk for transmission. Normal infection control measures should be in place, and staff should use respirators (as coughing in the patient can be accidentally induced by the procedure), eye protection, and non-sterile gloves.

Procedures for gastric aspiration adapted from WHO guidelines, 2006. An instructive video on the procedure can be found at the following website: https://www.youtube.com/watch?v=IWI_TY_LbZk&feature=youtu.be

Contraindications

- Child not fasted for 4 hours
- Low platelet count or bleeding tendency

Material required

- Non-sterile gloves
- Nasogastric tube (10F)
- Syringe 5-30 cc with appropriate connector for the nasogastric tube
- Litmus paper
- Specimen container
- Lab request forms
- Pen
- Sterile water or normal saline
- Sodium bicarbonate solution (8%)
- Alcohol/chlorhexidine

Procedure

- Position child on his/her back or side.
- Have an assistant to hold the child.
- Measure the distance between the nose and stomach to estimate the distance that will be required to insert the tube into the stomach.
- Attach a syringe to the nasogastric tube.
- Gently insert the nasogastric tube through the nose and advance it into the stomach.

- Withdraw gastric contents (2-5 mL) using the syringe attached to the nasogastric tube.
- To check that the position of the tube is correct, test the gastric contents with litmus paper: blue litmus turns red in response to acidic stomach contents. Tube position can also be checked by pushing 3-5 mL of air into the stomach and listening with a stethoscope over the stomach.
- If no fluid is aspirated, insert 5-10 mL of sterile water or normal saline and attempt to aspirate again. If still unsuccessful, attempt this again. Do not repeat more than three times.
- Withdraw gastric contents (ideally at least 5-10 mL).
- Transfer gastric fluid from the syringe into a sterile container.
- Add an equal volume of sodium bicarbonate to the specimen in order to neutralize the acidic gastric contents and prevent destruction of tubercle bacilli.

After the procedure

- Wipe the specimen container with alcohol/chlorhexidine to prevent cross- infection and label the container.
- Fill out the lab request forms.
- Transport the specimen in a cool box to the lab for processing as soon as possible (within 4 hours).
- Give the child his or her usual food.

Sputum Induction

Sputum induction (SI) is a useful procedure for obtaining sputum specimens in situations where suspected or known TB patients cannot self-expectorate, and where a bacteriological result is desired for diagnosis or follow up.

Practice points

- The procedure can be repeated twice on the same day, at least 4 hours apart, in order to obtain the specimens.
- Due to the risk of bronchospasm, only trained health staff must conduct the procedure, preferably a nurse.
- Sputum induction is an aerosol-generating procedure. Therefore, appropriate infection control measures must be taken. Specifically:
 - An appropriate site must be available. The minimum requirement is a small room with good ventilation.
 - Staff must use respirators, eye protection and non sterile gloves.

Material required

General

- Mask (respirator) for the operator and caregiver (if present)
- Eye protection and non sterile gloves for operator
- Oxygen (on standby in case of emergency)
- Pulse oximeter
- Request form

Preparation Pre-nebulization

- Spacer device (holding chamber) and mask
- Salbutamol metered dose inhaler

Nebulization

- Mask, chamber and tubing
- Antibacterial filter
- Nebulizer (Ultrasonic is the preferred type)
- Sterile solution of 3-6% sodium chloride, refrigerated if possible (more irritant)

Aspiration

- Suction material usually required only for children under 5 years old.
- Suction catheter (7 or 8F)
- Mechanical suction device & mucus trap or 50 mL syringe if not available
- Sputum collection container
- Sterile solution of 0.9% sodium chloride

Infection control measures

Management of materials

- Spacer devices (holding chambers) should either be sterilized after each patient (preferred) or disinfected after each patient by soaking in hexanios for at least 15 minutes, then rinse, then soak again in a new bath of hexanios for 15 minutes. Rinse well and then wipe dry.
- All masks, tubing, suction catheters and syringes should be disinfected with 2% chlorine and then discarded.
- Antibacterial filters should be fitted and changed for each patient to protect the nebulizer, oxygen cylinder (if used), and any aspiration device (if used).

Management of the environment

The site must be left unused with the windows open or extraction fan on for at least 30 minutes after the procedure to allow adequate replacement of air in the room. No one should enter this room during the period without a respirator.

Contraindications

- Patient not fasted for 2 hours
- Severe respiratory distress
- Oxygen saturation less than 92% in room air
- Bleeding – low platelet count, nose bleeds or other bleeding source
- Reduced level of consciousness
- History of significant asthma or chronic obstructive airways disease

Procedure

Prior to nebulization

- Explain the procedure to the patient and the accompanying adult.
- Have the patient in a sitting position.
- Ask older children to rinse their mouth with water.
- Use pulse oximeter to obtain baseline oxygen saturation.
- Administer 2 puffs of salbutamol 10 seconds apart. Use a holding chamber for all children. Wait 5 minutes before starting nebulization.
- Prepare a sputum container.

Nebulization

- Fill the nebulizer with 5 mL 3-6% hypertonic saline solution.
- Put on an N95 or FFP2 respirator and provide one for any accompanying adult.
- Place the nebulizer mask over the patient's face.
- Leave the patient to inhale.
- Stop the procedure and obtain a sample as soon as the patient starts to cough productively. In young children careful attention, with suctioning at the right moment is critical to avoid the sample being swallowed. If sputum is not induced during the procedure, continue until the reservoir is empty (not longer than 15 minutes), then attempt sample collection.

The patient should be observed for respiratory distress and the procedure should be stopped at any time if severe cough or wheeze develops.

Nasopharyngeal suction (usually required for children < 5 years)

- Do 1 to 2 minutes of clapping on the chest.
- Lay the child flat on his or her side, facing away from the operator.
- If a mechanical suction device and mucus extractor are available, use these. If not:
 - Fit a suction catheter to a 50 mL syringe. Lubricate the end of the catheter.
 - Measure the distance from the tip of the nose to the tragus of the ear.
 - Insert the suction catheter to that depth.
 - When inserting and withdrawing the tube, pull on the plunger of the syringe to create suction.
 - Once the syringe is filled with air and mucus, disconnect it from the suction catheter and purge the air (tip facing upward), so that only mucus is left in the syringe.
 - To collect the mucus, draw 2 mL of 0.9% saline into the syringe to rinse, then empty contents into the sample container.

Note that sputum may sometimes not be produced until up to 24 hours later. Therefore if a good sputum sample is not immediately produced, older children can be given a collection container to take home.

All patients should be observed for at least 15 minutes after the procedure to ensure there are no signs of respiratory distress. Recheck the oxygen saturation post procedure. Give oxygen if saturation has dropped below 90%.

Possible adverse effects to anticipate

In all cases, try to obtain a specimen only if the patient condition permits. Do not repeat the procedure in the case of severe adverse effects.

- Coughing spells (~40%)

If severe, stop the procedure and administer salbutamol. Oxygen should be available and can be administered in severe cases.

- Nosebleeds (~8%)

Stop the procedure and apply constant pressure to the mid portion of the nose until the bleeding stops. Note that it is very common to see blood in the specimens collected from nasopharyngeal suction; this in itself is not an adverse effect.

- Wheezing (<1%)

Monitor the child closely. Stop the procedure if wheeze increases. Administer salbutamol, and oxygen if severe.

- Vomiting (<1%)

Stop the procedure and observe the child closely until the vomiting stops.

Protocols courtesy of Michael Rich



The Sentinel Project on Pediatric Drug-Resistant Tuberculosis c/o the
Department of Global Health and Social Medicine Harvard Medical School

641 Huntington Avenue
Boston, MA 02115 USA

www.sentinel-project.org